

REGISTERED No. M - 302
L.-7646



EXTRAORDINARY
PUBLISHED BY AUTHORITY

ISLAMABAD, FRIDAY, DECEMBER, 31, 2021

PART III

Other Notifications, Orders, etc.

GOVERNMENT OF PAKISTAN

MINISTRY OF NATIONAL HEALTH SERVICES, REGULATIONS AND
COORDINATION

NOTIFICATION

Islamabad, the 21st December, 2021

No. F. 8-43/2021-DD(PS).—For the purposes of clause (xi) of subparagraph (1) of paragraph 2 of the Drug Pricing Policy, 2018, the Ministry of National Health Services, Regulations and Coordination is hereby pleased to notify the Model List of Essential Medicines, 2021 (22nd List) published by the World Health Organization.

Explanatory Notes:

The core list presents a list of minimum medicine needs for a basic health-care system, listing the most efficacious, safe and cost-effective medicines for priority conditions. Priority conditions are selected on the basis of current and estimated future public health relevance, and potential for safe and cost-effective treatment.

2131 (1—65)

Price: Rs. 100.00

[1918(2021)/Ex. Gaz.]

Where the [c] symbol is placed next to an individual medicine or strength of medicine on the core list it signifies that there is a specific indication for restricting its use to children.

The complementary list presents essential medicines for priority diseases, for which specialized diagnostic or monitoring facilities, and/or specialist medical care, and/or specialist training are needed. In case of doubt medicines may also be listed as complementary on the basis of consistent higher costs or less attractive cost-effectiveness in a variety of settings.

Where the [c] symbol is placed next to an individual medicine or strength of medicine on the complementary list it signifies that the medicine(s) require(s) specialist diagnostic or monitoring facilities, and/or specialist medical care, and/or specialist training for their use in children.

The square box symbol (□) is intended to indicate therapeutic alternatives to the listed medicine that may be considered for selection in national essential medicines lists. Alternatives may be individual medicines, or multiple medicines within a pharmacological class or chemical subgroup, defined at the 4th level of the Anatomical Therapeutic Chemical (ATC) classification, which have similar clinical effectiveness and safety. The listed medicine should be the example of the class or subgroup for which there is the best evidence for effectiveness and safety. In some cases, this may be the first medicine that is licensed for marketing; in other instances, subsequently licensed compounds may be safer or more effective. Where there is no difference in terms of efficacy and safety data, the listed medicine should be the one that is generally available at the lowest price, based on international drug price information sources. Not all square box listings are applicable to medicine selection for children. A square box is not used to indicate alternative generic brands of the same small molecule medicines, nor alternative biosimilars of biological medicines. However, the selection and use of quality-assured generics and biosimilars of essential medicines at country level is recommended.

National lists should not use a similar symbol and should be specific in their final selection, which would depend on local availability and price.

The [a] symbol indicates that there is an age or weight restriction on use of the medicine; details for each medicine can be found in Table 1.1.

The presence of an entry on the Essential Medicines List carries no assurance as to pharmaceutical quality. It is the responsibility of the relevant national or regional drug regulatory authority to ensure that each product is of

appropriate pharmaceutical quality (including stability) and that, when relevant, different products are interchangeable.

For recommendations and advice concerning all aspects of the quality assurance of medicines see the WHO website <https://www.who.int/teams/health-product-and-policy-standards/standards-and-specifications/norms-and-standards-for-pharmaceuticals/guidelines/quality-assurance>

Medicines and dosage forms are listed in alphabetical order within each section and the order of listing does not imply preference for one form over another. Standard treatment guidelines should be consulted for information on appropriate dosage forms.

The main terms used for dosage forms in the Essential Medicines List can be found in Table 1.2.

Definitions of many of these terms and pharmaceutical quality requirements applicable to the different categories are published in the current edition of *The International Pharmacopoeia*. <https://www.who.int/teams/health-product-and-policy-standards/standards-and-specifications/norms-and-standards-for-pharmaceuticals/pharmacopoeia>.

WHO Model List of Essential Medicines – 22nd List (2021)

ANAESTHETICS, PREOPERATIVE MEDICINES AND MEDICAL GASES	
1.1 General anaesthetics and oxygen	
1.1.1 Inhalational medicines	
halothane	Inhalation.
isoflurane	Inhalation.
nitrous oxide	Inhalation.
oxygen	Inhalation (medical gas).
1.1.2 Injectable medicines	
ketamine	Injection: 50 mg/mL (as hydrochloride) in 10 mL vial.
☐ propofol Therapeutic alternatives: - thiopental	Injection: 10 mg/mL; 20 mg/mL.
1.2 Local anaesthetics	
☐ bupivacaine Therapeutic alternatives to be reviewed (2023)	Injection: 0.25%; 0.5% (hydrochloride) in vial. Injection for spinal anaesthesia: 0.5% (hydrochloride) in 4 mL ampoule to be mixed with 7.5% glucose solution.
☐ lidocaine Therapeutic alternatives to be reviewed (2023)	Injection: 1%; 2% (hydrochloride) in vial. Injection for spinal anaesthesia: 5% (hydrochloride) in 2 mL ampoule to be mixed with 7.5% glucose solution. Topical forms: 2% to 4% (hydrochloride).
lidocaine + epinephrine (adrenaline)	Dental cartridge: 2% (hydrochloride) + epinephrine 1:80 000. Injection: 1%; 2% (hydrochloride or sulfate) + epinephrine 1:200 000 in vial.
<i>Complementary List</i>	
<i>ephedrine</i>	<i>Injection: 30 mg/mL (hydrochloride) in 1 mL ampoule. (For use in spinal anaesthesia during delivery, to prevent hypotension).</i>
1.3 Preoperative medication and sedation for short-term procedures	
atropine	Injection: 1 mg (sulfate) in 1 mL ampoule.
☐ midazolam Therapeutic alternatives to be reviewed (2023)	Injection: 1 mg/mL. Oral liquid: 2 mg/mL [c]. Tablet: 7.5 mg; 15 mg.
morphine	Injection: 10 mg (sulfate or hydrochloride) in 1 mL ampoule.

WHO Model List of Essential Medicines – 22nd List (2021)

1.4 Medical gases	
oxygen*	Inhalation For use in the management of hypoxaemia. *No more than 30% oxygen should be used to initiate resuscitation of neonates less than or equal to 32 weeks of gestation.
2. MEDICINES FOR PAIN AND PALLIATIVE CARE	
2.1 Non-opioids and non-steroidal anti-inflammatory medicines (NSAIDs)	
acetylsalicylic acid	Suppository: 50 mg to 150 mg. Tablet: 100 mg to 500 mg.
ibuprofen □	Oral liquid: 200 mg/5 mL. Tablet: 200 mg; 400 mg; 600 mg. □ Not in children less than 3 months.
paracetamol*	Oral liquid: 120 mg/5 mL; 125 mg/5 mL. Suppository: 100 mg. Tablet: 100 mg to 500 mg. *Not recommended for anti-inflammatory use due to lack of proven benefit to that effect.
2.2 Opioid analgesics	
codeine	Tablet: 30 mg (phosphate).
fentanyl*	Transdermal patch: 12 micrograms/hr; 25 micrograms/hr; 50 micrograms/hr; 75 micrograms/hr; 100 micrograms/hr *For the management of cancer pain
□ morphine Therapeutic alternatives: - hydromorphone - oxycodone	Granules (slow release; to mix with water): 20 mg to 200 mg (morphine sulfate). Injection: 10 mg (morphine hydrochloride or morphine sulfate) in 1 mL ampoule. Oral liquid: Tablet (slow release): 10 mg to 200 mg (morphine hydrochloride or morphine sulfate). Tablet (immediate release): 10 mg (morphine sulfate).
Complementary list	
methadone*	Tablet: 5 mg; 10 mg (hydrochloride) Oral liquid: 5 mg/5 mL; 10 mg/5 mL (hydrochloride) Concentrate for oral liquid: 5 mg/mL; 10 mg/mL (hydrochloride) *For the management of cancer pain.

WHO Model List of Essential Medicines – 22nd List (2021)

2.3 Medicines for other common symptoms in palliative care	
amitriptyline	Tablet: 10 mg; 25 mg; 75 mg.
cyclizine [c]	Injection: 50 mg/mL. Tablet: 50 mg.
dexamethasone	Injection: 4 mg/mL (as disodium phosphate salt) in 1 mL ampoule. Oral liquid: 2 mg/5 mL. Tablet: 2 mg [c]; 4 mg.
diazepam	Injection: 5 mg/mL. Oral liquid: 2 mg/5 mL. Rectal solution: 2.5 mg; 5 mg; 10 mg. Tablet: 5 mg; 10 mg.
docusate sodium	Capsule: 100 mg. Oral liquid: 50 mg/5 mL.
fluoxetine [a]	Solid oral dosage form: 20 mg (as hydrochloride). [a] > 6 years.
haloperidol	Injection: 5 mg in 1 mL ampoule. Oral liquid: 2 mg/mL. Solid oral dosage form: 0.5 mg; 2mg; 5 mg.
hyoscine butylbromide	Injection: 20 mg/mL.
hyoscine hydrobromide [c]	Injection: 400 micrograms/mL; 600 micrograms/mL. Transdermal patches: 1 mg/72 hours.
lactulose [c]	Oral liquid: 3.1 to 3.7 g/5 mL.
loperamide	Solid oral dosage form: 2 mg.
metoclopramide	Injection: 5 mg/mL (hydrochloride) in 2 mL ampoule. Oral liquid: 5 mg/5 mL. Solid oral form: 10 mg (hydrochloride).
midazolam	Injection: 1 mg/mL; 5 mg/mL. Oral liquid: 2mg/mL [c]. Solid oral dosage form: 7.5 mg; 15 mg.
□ ondansetron [a] Therapeutic alternatives: - dolasetron - granisetron - palonosetron - tropisetron	Injection: 2 mg base/mL in 2 mL ampoule (as hydrochloride). Oral liquid: 4 mg base/5 mL. Solid oral dosage form: Eq 4 mg base; Eq 8 mg base. [a] > 1 month.
senna	Oral liquid: 7.5 mg/5 mL.

WHO Model List of Essential Medicines – 22nd List (2021)

3. ANTIALLERGICS AND MEDICINES USED IN ANAPHYLAXIS	
dexamethasone	Injection: 4 mg/mL (as disodium phosphate salt) in 1 mL ampoule.
epinephrine (adrenaline)	Injection: 1 mg/mL (as hydrochloride or hydrogen tartrate) in 1 mL ampoule.
hydrocortisone	Powder for injection: 100 mg (as sodium succinate) in vial.
☐ loratadine* Therapeutic alternatives: - cetirizine - fexofenadine	Oral liquid: 1 mg/mL. Tablet: 10 mg. <i>*There may be a role for sedating antihistamines for limited indications (EMLc).</i>
☐ prednisolone Therapeutic alternatives: - prednisone	Oral liquid: 5 mg/mL [c]. Tablet: 5 mg; 25 mg.
4. ANTIDOTES AND OTHER SUBSTANCES USED IN POISONINGS	
4.1 Non-specific	
charcoal, activated	Powder.
4.2 Specific	
acetylcysteine	Injection: 200 mg/mL in 10 mL ampoule. Oral liquid: 10% [c]; 20% [c].
atropine	Injection: 1 mg (sulfate) in 1 mL ampoule.
calcium gluconate	Injection: 100 mg/mL in 10 mL ampoule.
methylthioninium chloride (methylene blue)	Injection: 10 mg/mL in 10 mL ampoule.
naloxone	Injection: 400 micrograms (hydrochloride) in 1 mL ampoule.
penicillamine	Solid oral dosage form: 250 mg.
potassium ferric hexacyano-ferrate(II) · 2H ₂ O (Prussian blue)	Powder for oral administration.
sodium nitrite	Injection: 30 mg/mL in 10 mL ampoule.
sodium thiosulfate	Injection: 250 mg/mL in 50 mL ampoule.
Complementary List	
deferoxamine	Powder for injection: 500 mg (mesilate) in vial.
dimercaprol	Injection in oil: 50 mg/mL in 2 mL ampoule.
fomepizole	Injection: 5 mg/mL (sulfate) in 20 mL ampoule or 1 g/mL (base) in 1.5 mL ampoule.
sodium calcium edetate	Injection: 200 mg/mL in 5 mL ampoule.
succimer	Solid oral dosage form: 100 mg.

WHO Model List of Essential Medicines – 22nd List (2021)

5. ANTICONVULSANTS/ANTIEPILEPTICS	
carbamazepine	Oral liquid: 100 mg/5 mL. Tablet (chewable): 100 mg; 200 mg. Tablet (scored): 100 mg; 200 mg.
diazepam	Gel or rectal solution: 5 mg/mL in 0.5 mL; 2 mL; 4 mL tubes.
lamotrigine*	Tablet: 25 mg; 50 mg; 100 mg; 200 mg. Tablet (chewable, dispersible): 2 mg; 5 mg; 25 mg; 50 mg; 100 mg; 200 mg. *For use as adjunctive therapy for treatment-resistant partial or generalized seizures.
☐ lorazepam Therapeutic alternatives: - diazepam (injection) - midazolam (injection)	Injection: 2 mg/mL in 1 mL ampoule; 4 mg/mL in 1 mL ampoule.
magnesium sulfate*	Injection: 0.5 g/mL in 2 mL ampoule (equivalent to 1 g in 2 mL; 50% weight/volume); 0.5 g/mL in 10 mL ampoule (equivalent to 5 g in 10 mL; 50% weight/volume). *For use in eclampsia and severe pre-eclampsia and not for other convulsant disorders.
midazolam	Solution for oromucosal administration: 5 mg/mL; 10 mg/mL. Ampoule*: 1 mg/mL; 10 mg/mL. *For buccal administration when solution for oromucosal administration is not available.
phenobarbital	Injection: 200 mg/mL (sodium). Oral liquid: 15 mg/5 mL. Tablet: 15 mg to 100 mg.
phenytoin	Injection: 50 mg/mL (sodium) in 5 mL vial. Oral liquid: 25 mg to 30 mg/5 mL.* Solid oral dosage form: 25 mg; 50 mg; 100 mg (sodium). Tablet (chewable): 50 mg. *The presence of both 25 mg/5 mL and 30 mg/5 mL strengths on the same market would cause confusion in prescribing and dispensing and should be avoided.
valproic acid (sodium valproate)* <i>*Avoid use in pregnancy and in women and girls of child-bearing potential, unless alternative treatments are ineffective or not tolerated because of the high risk of birth defects and developmental disorders in children exposed to valproate in the womb.</i>	Oral liquid: 200 mg/5 mL. Tablet (crushable): 100 mg. Tablet (enteric-coated): 200 mg; 500 mg.

WHO Model List of Essential Medicines – 22nd List (2021)

<i>Complementary List</i>	
ethosuximide	Capsule: 250 mg. Oral liquid: 250 mg/5 mL.
valproic acid (sodium valproate)* <i>*Avoid use in pregnancy and in women and girls of child-bearing potential, unless alternative treatments are ineffective or not tolerated because of the high risk of birth defects and developmental disorders in children exposed to valproate in the womb.</i>	Injection: 100 mg/mL in 4 mL ampoule; 100 mg/mL in 10 mL ampoule.
6. ANTI-INFECTIVE MEDICINES	
6.1 Anthelmintics	
6.1.1 Intestinal anthelmintics	
albendazole	Tablet (chewable): 400 mg.
ivermectin	Tablet (scored): 3 mg.
levamisole	Tablet: 50 mg; 150 mg (as hydrochloride).
mebendazole	Tablet (chewable): 100 mg; 500 mg.
niclosamide	Tablet (chewable): 500 mg.
praziquantel	Tablet: 150 mg; 600 mg.
pyrantel	Oral liquid: 50 mg/mL (as embonate or pamoate). Tablet (chewable): 250 mg (as embonate or pamoate).
6.1.2 Antifilarials	
albendazole	Tablet (chewable): 400 mg.
diethylcarbamazine	Tablet: 50 mg; 100 mg (dihydrogen citrate).
ivermectin	Tablet (scored): 3 mg.
6.1.3 Antischistosomes and other antitrepanematode medicines	
praziquantel	Tablet: 600 mg.
triclabendazole	Tablet: 250 mg.
<i>Complementary List</i>	
oxamniquine*	Capsule: 250 mg. Oral liquid: 250 mg/5 mL. <i>*For use when praziquantel treatment fails.</i>
6.1.4 Cysticidal medicines	
<i>Complementary List</i>	
albendazole	Tablet (chewable): 400 mg.
mebendazole	Tablet (chewable): 500 mg.
praziquantel	Tablet: 500 mg; 600 mg.

WHO Model List of Essential Medicines – 22nd List (2021)

6.2 Antibacterials

To assist in the development of tools for antibiotic stewardship at local, national and global levels and to reduce antimicrobial resistance, the Access, Watch, Reserve (AWaRe) classification of antibiotics was developed – where antibiotics are classified into different groups to emphasize the importance of their appropriate use.

ACCESS GROUP ANTIBIOTICS

This group includes antibiotics that have activity against a wide range of commonly encountered susceptible pathogens while also showing lower resistance potential than antibiotics in the other groups. Selected Access group antibiotics are recommended as essential first or second choice empiric treatment options for infectious syndromes reviewed by the EML Expert Committee and are listed as individual medicines on the Model Lists to improve access and promote appropriate use. They are essential antibiotics that should be widely available, affordable and quality assured.

WATCH GROUP ANTIBIOTICS

This group includes antibiotic classes that have higher resistance potential and includes most of the highest priority agents among the Critically Important Antimicrobials for Human Medicine and/or antibiotics that are at relatively high risk of selection of bacterial resistance. These medicines should be prioritized as key targets of stewardship programs and monitoring. Selected Watch group antibiotics are recommended as essential first or second choice empiric treatment options for a limited number of specific infectious syndromes and are listed as individual medicines on the Model Lists.

RESERVE GROUP ANTIBIOTICS

This group includes antibiotics and antibiotic classes that should be reserved for treatment of confirmed or suspected infections due to multi-drug-resistant organisms. Reserve group antibiotics should be treated as "last resort" options. Selected Reserve group antibiotics are listed as individual medicines on the Model Lists when they have a favourable risk-benefit profile and proven activity against "Critical Priority" or "High Priority" pathogens identified by the WHO Priority Pathogens List, notably carbapenem resistant *Enterobacteriaceae*. These antibiotics should be accessible, but their use should be tailored to highly specific patients and settings, when all alternatives have failed or are not suitable. These medicines could be protected and prioritized as key targets of national and international stewardship programs involving monitoring and utilization reporting, to preserve their effectiveness.

WHO Model List of Essential Medicines – 22nd List (2021)

6.2.1 Access group antibiotics		
amikacin	Injection: 250 mg/mL (as sulfate) in 2 mL vial.	
	FIRST CHOICE – High-risk febrile neutropenia – Pyelonephritis or prostatitis (severe)	SECOND CHOICE – Sepsis in neonates and children [c]
amoxicillin	Powder for injection: 250 mg; 500 mg; 1 g (as sodium) in vial. Powder for oral liquid: 125 mg/5 mL; 250 mg/5 mL (as trihydrate) [c]. Solid oral dosage form: 250 mg; 500 mg; 1g (as trihydrate).	
	FIRST CHOICE – Community acquired pneumonia (mild to moderate) – Community acquired pneumonia (severe) [c] – Complicated severe acute malnutrition [c] – Exacerbations of COPD – Otitis media – Pharyngitis – Progressive apical dental abscess – Sepsis in neonates and children [c] – Sinusitis – Uncomplicated severe acute malnutrition [c]	SECOND CHOICE – Acute bacterial meningitis
amoxicillin + clavulanic acid	Powder for injection: 500 mg (as sodium) + 100 mg (as potassium salt); 1000 mg (as sodium) + 200 mg (as potassium salt) in vial. Powder for oral liquid: 125 mg (as trihydrate) + 31.25 mg (as potassium salt)/5 mL; 250 mg (as trihydrate) + 62.5 mg (as potassium salt)/5 mL [c]. Tablet: 500 mg (as trihydrate) + 125 mg (as potassium salt); 875 mg (as trihydrate) + 125 mg (as potassium salt).	
	FIRST CHOICE – Community acquired pneumonia (severe) [c] – Complicated intraabdominal infections (mild to moderate) – Exacerbations of COPD – Hospital acquired pneumonia – Low-risk febrile neutropenia – Lower urinary tract infections – Sinusitis – Skin and soft tissue infections	SECOND CHOICE – Bone and joint infections – Community-acquired pneumonia (mild to moderate) – Community acquired pneumonia (severe) – Otitis media – Surgical prophylaxis

WHO Model List of Essential Medicines – 22nd List (2021)

ampicillin	Powder for injection: 500 mg; 1 g (as sodium) in vial.	
	FIRST CHOICE – Community acquired pneumonia (severe) [c] – Complicated intraabdominal infections [c] – Complicated severe acute malnutrition [c] – Sepsis in neonates and children [c]	SECOND CHOICE – Acute bacterial meningitis
benzathine benzylpenicillin	Powder for injection: 1.2 million IU (= 900 mg) in vial [c]; 2.4 million IU (= 1.8 g) in vial.	
	FIRST CHOICE – Syphilis	SECOND CHOICE
benzylpenicillin	Powder for injection: 600 mg (= 1 million IU); 3 g (= 5 million IU) (sodium or potassium salt) in vial.	
	FIRST CHOICE – Community acquired pneumonia (severe) [c] – Complicated severe acute malnutrition [c] – Sepsis in neonates and children [c] – Syphilis	SECOND CHOICE – Acute bacterial meningitis [c]
cefalexin	Powder for oral liquid: 125 mg/5 mL; 250 mg/5 mL (anhydrous). Solid oral dosage form: 250 mg; 500 mg (as monohydrate).	
	FIRST CHOICE – Skin and soft tissue infections	SECOND CHOICE – Exacerbations of COPD – Pharyngitis
cefazolin ^a	Powder for injection: 1 g (as sodium salt) in vial. ^a > 1 month.	
	FIRST CHOICE – Surgical prophylaxis	SECOND CHOICE – Bone and joint infections
chloramphenicol	Capsule: 250 mg. Oily suspension for injection*: 0.5 g/mL (as sodium succinate) in 2 mL ampoule. *Only for the presumptive treatment of epidemic meningitis in children older than 2 years and in adults. Oral liquid: 150 mg/5 mL (as palmitate). Powder for injection: 1 g (sodium succinate) in vial.	
	FIRST CHOICE	SECOND CHOICE – Acute bacterial meningitis

WHO Model List of Essential Medicines – 22nd List (2021)

clindamycin	Capsule: 150 mg (as hydrochloride). Injection: 150 mg/mL (as phosphate); 600 mg/4 mL (as phosphate); 900 mg/6 mL (as phosphate). Oral liquid: 75 mg/5 mL (as palmitate) [c].	
	FIRST CHOICE – Necrotizing fasciitis	SECOND CHOICE – Bone and joint infections
□ cloxacillin* Therapeutic alternatives: – 4 th level ATC chemical subgroup (J01CF Beta-lactamase resistant penicillins)	Capsule: 500 mg; 1 g (as sodium). Powder for injection: 500 mg (as sodium) in vial. Powder for oral liquid: 125 mg/5 mL (as sodium). *cloxacillin, dicloxacillin and flucloxacillin are preferred for oral administration due to better bioavailability.	
	FIRST CHOICE – Bone and joint infections – Skin and soft tissue infections	SECOND CHOICE – Sepsis in neonates and children [c]
doxycycline ^a	Oral liquid: 25 mg/5 mL [c]; 50 mg/5 mL (anhydrous) [c]. Powder for injection: 100 mg in vial. Solid oral dosage form: 50 mg [c]; 100 mg (as hyclate). ^a Use in children <8 years only for life-threatening infections when no alternative exists.	
	FIRST CHOICE – Cholera – Sexually transmitted infection due to <i>Chlamydia trachomatis</i>	SECOND CHOICE – Cholera [c] – Community acquired pneumonia (mild to moderate) – Exacerbations of COPD
gentamicin	Injection: 10 mg/mL (as sulfate); 40 mg/mL (as sulfate) in 2 mL vial.	
	FIRST CHOICE – Acute bacterial meningitis in neonates [c] – Community acquired pneumonia (severe) [c] – Complicated intraabdominal infections [c] – Complicated severe acute malnutrition [c] – Sepsis in neonates and children [c]	SECOND CHOICE – Gonorrhoea – Surgical prophylaxis

WHO Model List of Essential Medicines – 22nd List (2021)

metronidazole	Injection: 500 mg in 100 mL vial. Oral liquid: 200 mg/5 mL (as benzoate). Suppository: 500 mg; 1 g. Tablet: 200 mg to 500 mg.	
	FIRST CHOICE – <i>C. difficile</i> infection – Complicated intraabdominal infections (mild to moderate) – Complicated intrabdominal infections (severe) – Necrotizing fasciitis – Surgical prophylaxis – Trichomoniasis	SECOND CHOICE – Complicated intraabdominal infections (mild to moderate)
nitrofurantoin	Oral liquid: 25 mg/5 mL [c]. Tablet: 100 mg.	
	FIRST CHOICE – Lower urinary tract infections	SECOND CHOICE
phenoxymethylpenicillin	Powder for oral liquid: 250 mg/5 mL (as potassium). Tablet: 250 mg; 500 mg (as potassium).	
	FIRST CHOICE – Community acquired pneumonia (mild to moderate) – Pharyngitis – Progressive apical dental abscess	SECOND CHOICE
procaine benzylpenicillin*	Powder for injection: 1 g (=1 million IU); 3 g (=3 million IU) in vial. *Procaine benzylpenicillin is not recommended as first-line treatment for neonatal sepsis except in settings with high neonatal mortality, when given by trained health workers in cases where hospital care is not achievable.	
	FIRST CHOICE – Syphilis (congenital) [c]	SECOND CHOICE – Syphilis
spectinomycin	Powder for injection: 2 g (as hydrochloride) in vial.	
	FIRST CHOICE	SECOND CHOICE – Gonorrhoea
sulfamethoxazole + trimethoprim	Injection: 80 mg + 16 mg/mL in 5 mL ampoule; 80 mg + 16 mg/mL in 10 mL ampoule. Oral liquid: 200 mg + 40 mg/5 mL. Tablet: 100 mg + 20 mg; 400 mg + 80 mg; 800 mg + 160 mg.	
	FIRST CHOICE – Lower urinary tract infections	SECOND CHOICE – Acute invasive diarrhoea/bacterial dysentery

WHO Model List of Essential Medicines – 22nd List (2021)

trimethoprim	Tablet: 100 mg; 200 mg. Oral liquid: 50 mg/5 mL [c].	
	FIRST CHOICE – Lower urinary tract infections	SECOND CHOICE
6.2.2 Watch group antibiotics		
azithromycin	Capsule: 250 mg; 500 mg (anhydrous). Oral liquid: 200 mg/5 mL.	
	FIRST CHOICE – Cholera [c] – Enteric fever – Gonorrhoea – Sexually transmitted infection due to <i>Chlamydia trachomatis</i> – Trachoma – Yaws	SECOND CHOICE – Acute invasive bacterial diarrhoea / dysentery – Gonorrhoea
cefixime	Powder for oral liquid: 100 mg/5 mL [c]. Solid oral dosage form: 200 mg; 400 mg (as trihydrate).	
	FIRST CHOICE	SECOND CHOICE – Acute invasive bacterial diarrhoea / dysentery – Gonorrhoea
cefotaxime*	Powder for injection: 250 mg (as sodium) in vial. *3rd generation cephalosporin of choice for use in hospitalized neonates.	
	FIRST CHOICE – Acute bacterial meningitis – Community acquired pneumonia (severe) – Complicated intraabdominal infections (mild to moderate) – Complicated intraabdominal infections (severe) – Hospital acquired pneumonia – Pyelonephritis or prostatitis (severe)	SECOND CHOICE – Bone and joint infections – Pyelonephritis or prostatitis (mild to moderate) – Sepsis in neonates and children [c]

WHO Model List of Essential Medicines – 22nd List (2021)

ceftriaxone*†§	Powder for injection: 250 mg; 1 g; 2 g (as sodium) in vial. *Do not administer with calcium and avoid in infants with hyperbilirubinaemia. § > 41 weeks corrected gestational age.	
	FIRST CHOICE – Acute bacterial meningitis – Community acquired pneumonia (severe) – Complicated intraabdominal infections (mild to moderate) – Complicated intraabdominal infections (severe) – Endophthalmitis – Enteric fever – Gonorrhoea – Hospital acquired pneumonia – Necrotizing fasciitis – Pyelonephritis or prostatitis (severe)	SECOND CHOICE – Acute invasive bacterial diarrhoea / dysentery – Bone and joint infections – Pyelonephritis or prostatitis (mild to moderate) – Sepsis in neonates and children[c]
cefuroxime	Powder for injection: 250 mg; 750 mg; 1.5 g (as sodium) in vial.	
	FIRST CHOICE	SECOND CHOICE – Surgical prophylaxis
ciprofloxacin	Oral liquid: 250 mg/5 mL (anhydrous) [c]. Solution for IV infusion: 2 mg/mL (as hyclate) [c]. Solid oral dosage form: 250 mg; 500 mg (as hydrochloride).	
	FIRST CHOICE – Acute invasive bacterial diarrhoea / dysentery – Enteric fever – Low-risk febrile neutropenia – Pyelonephritis or prostatitis (mild to moderate)	SECOND CHOICE – Cholera – Complicated intraabdominal infections (mild to moderate)
□ clarithromycin† Therapeutic alternatives: – erythromycin* *as second choice treatment for pharyngitis in children (EMLc only)	Powder for oral liquid: 125 mg/5 mL; 250 mg/5 mL. Powder for injection: 500 mg in vial. Solid oral dosage form: 500 mg. †clarithromycin is also listed for use in combination regimens for eradication of <i>H. pylori</i> in adults.	
	FIRST CHOICE – Community acquired pneumonia (severe)	SECOND CHOICE – Pharyngitis

WHO Model List of Essential Medicines – 22nd List (2021)

piperacillin + tazobactam	Powder for injection: 2 g (as sodium) + 250 mg (as sodium); 4 g (as sodium) + 500 mg (as sodium) in vial.	
	FIRST CHOICE – Complicated intraabdominal infections (severe) – High-risk febrile neutropenia – Hospital acquired pneumonia – Necrotizing fasciitis	SECOND CHOICE
vancomycin	Capsule: 125 mg; 250 mg (as hydrochloride).	
	FIRST CHOICE	SECOND CHOICE – <i>C. difficile</i> infection
Complementary List		
ceftazidime	Powder for injection: 250 mg; 1 g (as pentahydrate) in vial.	
	FIRST CHOICE – Endophthalmitis	SECOND CHOICE
☐ meropenem ^a Therapeutic alternatives ^b : – imipenem + cilastatin ^a > 3 months. ^b complicated intraabdominal infections and high-risk febrile neutropenia only. Meropenem is the preferred choice for acute bacterial meningitis in neonates.	Powder for injection: 500 mg (as trihydrate); 1 g (as trihydrate) in vial.	
	FIRST CHOICE	SECOND CHOICE – Acute bacterial meningitis in neonates [c] – Complicated intraabdominal infections (severe) – High-risk febrile neutropenia
vancomycin	Powder for injection: 250 mg; 500 mg; 1 g (as hydrochloride) in vial.	
	FIRST CHOICE – Endophthalmitis – Necrotizing fasciitis	SECOND CHOICE – High-risk febrile neutropenia
6.2.3 Reserve group antibiotics		
Complementary List		
cefiderocol	Powder for injection: 1 g (as sulfate toxyrate) in vial.	
ceftazidime + avibactam	Powder for injection: 2 g + 0.5 g in vial.	
colistin	Powder for injection: 1 million IU (as colistimethate sodium) in vial.	
fosfomycin	Powder for injection: 2 g; 4 g (as sodium) in vial.	
linezolid	Injection for intravenous administration: 2 mg/mL in 300 mL bag. Powder for oral liquid: 100 mg/5 mL. Tablet: 400 mg; 600 mg.	
meropenem + vaborbactam	Powder for injection: 1 g (as trihydrate) + 1 g in vial.	
plazomicin	Injection: 500 mg/10 mL.	
polymyxin B	Powder for injection: 500,000 IU in vial.	

WHO Model List of Essential Medicines – 22nd List (2021)

6.2.4 Antileprosy medicines	
Medicines used in the treatment of leprosy should never be used except in combination. Combination therapy is essential to prevent the emergence of drug resistance. Colour-coded blister packs (MDT blister packs) containing standard two-medicine (paucibacillary leprosy) or three-medicine (multibacillary leprosy) combinations for adult and childhood leprosy should be used. MDT blister packs can be supplied free of charge through WHO.	
clofazimine	Capsule: 50 mg; 100 mg.
dapsone	Tablet: 25 mg; 50 mg; 100 mg.
rifampicin	Solid oral dosage form: 150 mg; 300 mg.
6.2.5 Antituberculosis medicines	
WHO recommends and endorses the use of fixed-dose combinations and the development of appropriate new fixed-dose combinations, including modified dosage forms, non-refrigerated products and paediatric dosage forms of assured pharmaceutical quality.	
ethambutol	Oral liquid: 25 mg/mL [c]. Tablet: 100 mg; 400 mg (hydrochloride). Tablet (dispersible): 100 mg [c]
ethambutol + isoniazid + pyrazinamide + rifampicin	Tablet: 275 mg + 75 mg + 400 mg + 150 mg.
ethambutol + isoniazid + rifampicin	Tablet: 275 mg + 75 mg + 150 mg.
isoniazid	Oral liquid: 50 mg/5 mL [c]. Tablet: 100 mg; 300 mg. Tablet (dispersible): 100 mg [c].
isoniazid + pyrazinamide + rifampicin	Tablet (dispersible): 50 mg + 150 mg + 75 mg [c].
isoniazid + rifampicin	Tablet: 75 mg + 150 mg; 150 mg + 300 mg. Tablet (dispersible): 50 mg + 75 mg [c].
isoniazid + rifapentine	Tablet (scored): 300 mg + 300 mg.
moxifloxacin	Tablet: 400 mg.
pyrazinamide	Oral liquid: 30 mg/mL [c]. Tablet: 400 mg; 500 mg. Tablet (dispersible): 150 mg.
rifabutin	Solid oral dosage form: 150 mg.* *For use only in patients with HIV receiving protease inhibitors.
rifampicin	Oral liquid: 20 mg/mL [c]. Solid oral dosage form: 150 mg; 300 mg.
rifapentine	Tablet: 150 mg; 300 mg.

WHO Model List of Essential Medicines – 22nd List (2021)

Complementary List	
<i>Medicines for the treatment of multidrug-resistant tuberculosis (MDR-TB) should be used in specialized centres adhering to WHO standards for TB control.</i>	
amikacin	Injection: 100 mg/2 mL (as sulfate) in 2 mL vial; 250 mg/mL (as sulfate) in 2 mL vial.
amoxicillin + clavulanic acid*	Powder for oral liquid: 250 mg (as trihydrate) + 62.5 mg (as potassium salt)/5 mL [c]. Tablet: 500 mg (as trihydrate) + 125 mg (as potassium salt). *For use only in combination with meropenem or imipenem + cilastatin.
bedaquiline ^a	Tablet: 20 mg [c]; 100 mg. ^a ≥ 5 years
clofazimine	Solid oral dosage form: 50 mg; 100 mg.
☐ cycloserine Therapeutic alternatives: - terizidone	Solid oral dosage form: 125 mg [c]; 250 mg.
delamanid ^a	Tablet (dispersible): 25 mg [c]. ^a ≥ 3 years Tablet: 50 mg. ^a ≥ 6 years
☐ ethionamide Therapeutic alternatives: - protonamide	Tablet: 125 mg; 250 mg. Tablet (dispersible): 125 mg [c].
levofloxacin	Tablet: 250 mg; 500 mg; 750 mg. Tablet (dispersible): 100 mg [c].
linezolid	Powder for oral liquid: 100 mg/5 mL. Tablet: 600 mg. Tablet (dispersible): 150 mg [c].
☐ meropenem Therapeutic alternatives: - imipenem + cilastatin	Powder for injection: 500 mg (as trihydrate); 1 g (as trihydrate) in vial.
moxifloxacin	Tablet: 400 mg. Tablet (dispersible): 100 mg [c].
p-aminosalicylic acid	Granules: 4 g in sachet.
streptomycin [c]	Powder for injection: 1 g (as sulfate) in vial.

WHO Model List of Essential Medicines – 22nd List (2021)

6.3 Antifungal medicines	
amphotericin B	Powder for injection: 50 mg (as sodium deoxycholate or liposomal complex) in vial.
clotrimazole	Vaginal cream: 1%; 10%. Vaginal tablet: 100 mg; 500 mg.
fluconazole	Capsule: 50 mg. Injection: 2 mg/mL in vial. Oral liquid: 50 mg/5 mL.
flucytosine	Capsule: 250 mg. Infusion: 2.5 g in 250 mL.
griseofulvin	Oral liquid: 125 mg/5 mL [c]. Solid oral dosage form: 125 mg; 250 mg.
itraconazole*	Capsule: 100 mg. Oral liquid: 10 mg/mL. *For treatment of chronic pulmonary aspergillosis, histoplasmosis, sporotrichosis, paracoccidioidomycosis, mycoses caused by <i>T. marneffei</i> and chromoblastomycosis; and prophylaxis of histoplasmosis and infections caused by <i>T. marneffei</i> in AIDS patients.
nystatin	Lozenge: 100 000 IU. Oral liquid: 50 mg/5 mL [c]; 100 000 IU/mL [c]. Pessary: 100 000 IU. Tablet: 100 000 IU; 500 000 IU.
voriconazole*	Tablet: 50 mg; 200 mg Powder for injection: 200 mg in vial Powder for oral liquid: 40 mg/mL *For treatment of chronic pulmonary aspergillosis and acute invasive aspergillosis.
Complementary List	
□ micafungin Therapeutic alternatives: - anidulafungin - caspofungin	Powder for injection: 50 mg (as sodium); 100 mg (as sodium) in vial.
potassium iodide	Saturated solution.

WHO Model List of Essential Medicines – 22nd List (2021)

6.4 Antiviral medicines	
6.4.1 <i>Antiherpes medicines</i>	
☐ aciclovir Therapeutic alternatives: - valaciclovir (oral)	Oral liquid: 200 mg/5 mL [c]. Powder for injection: 250 mg (as sodium salt) in vial. Tablet: 200 mg.
6.4.2 <i>Antiretrovirals</i>	
Based on current evidence and experience of use, medicines in the following classes of antiretrovirals are included as essential medicines for treatment and prevention of HIV (prevention of mother-to-child transmission, pre-exposure prophylaxis (where indicated) and post-exposure prophylaxis). WHO emphasizes the importance of using these products in accordance with global and national guidelines. WHO recommends and endorses the use of fixed-dose combinations and the development of appropriate new fixed-dose combinations, including modified dosage forms, non-refrigerated products and paediatric dosage forms of assured pharmaceutical quality. Scored tablets can be used in children and therefore can be considered for inclusion in the listing of tablets, provided that adequate quality products are available.	
6.4.2.1 <i>Nucleoside/Nucleotide reverse transcriptase inhibitors</i>	
abacavir	Tablet: 300 mg (as sulfate).
lamivudine	Oral liquid: 50 mg/5 mL [c]. Tablet: 150 mg.
tenofovir disoproxil fumarate†	Tablet: 300 mg (tenofovir disoproxil fumarate – equivalent to 245 mg tenofovir disoproxil). †also indicated for pre-exposure prophylaxis.
zidovudine	Capsule: 250 mg. Oral liquid: 50 mg/5 mL. Solution for IV infusion: 10 mg/mL in 20 mL vial. Tablet: 300 mg.
6.4.2.2 <i>Non-nucleoside reverse transcriptase inhibitors</i>	
efavirenz	Tablet: 600 mg.
nevirapine ^a	Oral liquid: 50 mg/5 mL. Tablet: 50 mg (dispersible); 200 mg. ^a > 6 weeks
6.4.2.3 <i>Protease inhibitors</i>	
Selection of protease inhibitor(s) from the Model List will need to be determined by each country after consideration of international and national treatment guidelines and experience. Ritonavir is recommended for use in combination as a pharmacological booster, and not as an antiretroviral in its own right. All other protease inhibitors should be used in boosted forms (e.g. with ritonavir).	
atazanavir + ritonavir	Tablet (heat stable): 300 mg (as sulfate) + 100 mg.
darunavir ^a	Tablet: 75 mg; 400 mg; 600 mg; 800 mg ^a > 3 years
lopinavir + ritonavir	Solid oral dosage form: 40 mg + 10 mg [c]. Tablet (heat stable): 100 mg + 25 mg; 200 mg + 50 mg.
ritonavir	Tablet (heat stable): 25 mg; 100 mg.

WHO Model List of Essential Medicines – 22nd List (2021)

6.4.2.4 <i>Integrase inhibitors</i>	
dolutegravir ^a	Tablet (dispersible, scored): 10 mg [c]. ^a ≥ 4 weeks and ≥ 3 kg Tablet: 50 mg ^a ≥ 25 kg
raltegravir*	Granules for oral suspension: 100 mg in sachet. Tablet (chewable): 25 mg. Tablet: 400 mg. *For use in pregnant women and in second-line regimens in accordance with WHO treatment guidelines.
6.4.2.5 <i>Fixed-dose combinations of antiretroviral medicines</i>	
abacavir + lamivudine	Tablet (dispersible, scored): 120 mg (as sulfate) + 60 mg.
dolutegravir + lamivudine + tenofovir	Tablet: 50 mg + 300 mg + 300 mg (tenofovir disoproxil fumarate – equivalent to 245 mg tenofovir disoproxil)
efavirenz + □ emtricitabine + tenofovir Therapeutic alternatives: - lamivudine (for emtricitabine)	Tablet: 600 mg + 200 mg + 300 mg (tenofovir disoproxil fumarate – equivalent to 245 mg tenofovir disoproxil).
efavirenz + lamivudine + tenofovir	Tablet: 400 mg + 300 mg + 300 mg (tenofovir disoproxil fumarate – equivalent to 245 mg tenofovir disoproxil)
□ emtricitabine + tenofovir† Therapeutic alternatives: - lamivudine (for emtricitabine)	Tablet: 200 mg + 300 mg (tenofovir disoproxil fumarate – equivalent to 245 mg tenofovir disoproxil). † combination also indicated for pre-exposure prophylaxis
lamivudine + zidovudine	Tablet: 30 mg + 60 mg [c]; 150 mg + 300 mg.
6.4.2.6 <i>Medicines for prevention of HIV-related opportunistic infections</i>	
isoniazid + pyridoxine + sulfamethoxazole + trimethoprim	Tablet (scored): 300 mg + 25 mg + 800 mg + 160 mg
6.4.3 <i>Other antivirals</i>	
ribavirin*	Injection for intravenous administration: 800 mg and 1 g in 10 mL phosphate buffer solution. Solid oral dosage form: 200 mg; 400 mg; 600 mg. *For the treatment of viral haemorrhagic fevers
valganciclovir*	Tablet: 450 mg. *For the treatment of cytomegalovirus retinitis (CMVr).

WHO Model List of Essential Medicines – 22nd List (2021)

<i>Complementary list</i>	
oseltamivir*	Capsule: 30 mg; 45 mg; 75 mg (as phosphate). *Severe illness due to confirmed or suspected influenza virus infection in critically ill hospitalized patients
valganciclovir*[c]	Powder for oral solution: 50 mg/mL Tablet: 450 mg. *For the treatment of cytomegalovirus retinitis (CMVr).
6.4.4 Antihepatitis medicines	
6.4.4.1 Medicines for hepatitis B	
6.4.4.1.1 Nucleoside/Nucleotide reverse transcriptase inhibitors	
entecavir	Oral liquid: 0.05 mg/mL Tablet: 0.5 mg; 1 mg
tenofovir disoproxil fumarate	Tablet: 300 mg (tenofovir disoproxil fumarate – equivalent to 245 mg tenofovir disoproxil).
6.4.4.2 Medicines for hepatitis C	
Pangenotypic direct-acting antivirals should be considered as therapeutic alternatives for the purposes of selection and procurement at national level.	
6.4.4.2.1 ☐ Pangenotypic direct-acting antiviral combinations	
daclatasvir*	Tablet: 30 mg; 60 mg (as hydrochloride). *Pangenotypic when used in combination with sofosbuvir
daclatasvir + sofosbuvir	Tablet: 60 mg + 400 mg.
glecaprevir + pibrentasvir	Tablet: 100 mg + 40 mg. Granules: 50 mg + 20 mg in sachet [c].
sofosbuvir*	Tablet: 200 mg; 400 mg. *Pangenotypic when used in combination with daclatasvir
sofosbuvir + velpatasvir	Tablet: 200 mg + 50 mg [c]; 400 mg + 100 mg.
6.4.4.2.2 Non-pangenotypic direct-acting antiviral combinations	
dasabuvir	Tablet: 250 mg.
ledipasvir + sofosbuvir	Tablet: 90 mg + 400 mg.
ombitasvir + paritaprevir + ritonavir	Tablet: 12.5 mg + 75 mg + 50 mg.

WHO Model List of Essential Medicines – 22nd List (2021)

6.4.4.2.3 Other antivirals for hepatitis C	
ribavirin*	Injection for intravenous administration: 800 mg and 1 g in 10 mL phosphate buffer solution. Solid oral dosage form: 200 mg; 400 mg; 600 mg. *For the treatment of hepatitis C, in combination with direct acting anti-viral medicines
Complementary list	
pegylated interferon alfa (2a or 2b) *	Vial or pre-filled syringe: 180 micrograms (peginterferon alfa-2a). 80 micrograms, 100 micrograms (peginterferon alfa-2b). *To be used in combination with ribavirin.
6.5 Antiprotozoal medicines	
6.5.1 Antiamoebic and anti giardiasis medicines	
diloxanide [a]	Tablet: 500 mg (furoate). [a] > 25 kg.
metronidazole Therapeutic alternatives: - tinidazole	Injection: 500 mg in 100 mL vial. Oral liquid: 200 mg/5 mL (as benzoate). Tablet: 200 mg to 500 mg.
6.5.2 Antileishmaniasis medicines	
amphotericin B	Powder for injection: 50 mg in vial (as sodium deoxycholate or liposomal complex).
miltefosine	Solid oral dosage form: 10 mg; 50 mg.
paromomycin	Solution for intramuscular injection: 750 mg of paromomycin base (as sulfate).
sodium stibogluconate or meglumine antimoniate	Injection: 100 mg/mL, 1 vial = 30 mL or 30%, equivalent to approximately 8.1% antimony (pentavalent) in 5 mL ampoule.
6.5.3 Antimalarial medicines	
6.5.3.1 For curative treatment	
Medicines for the treatment of <i>P. falciparum</i> malaria cases should be used in combination. The list currently recommends combinations according to treatment guidelines. WHO recognizes that not all of the fixed dose combinations (FDCs) in the WHO treatment guidelines exist, and encourages their development and rigorous testing. WHO also encourages development and testing of rectal dosage formulations.	
amodiaquine*	Tablet: 153 mg or 200 mg (as hydrochloride). *To be used in combination with artesunate 50 mg.
artemether*	Oily injection: 80 mg/mL in 1 mL ampoule. *For use in the management of severe malaria.
artemether + lumefantrine*	Tablet: 20 mg + 120 mg. Tablet (dispersible): 20 mg + 120 mg (c). *Not recommended in the first trimester of pregnancy or in children below 5 kg.

WHO Model List of Essential Medicines – 22nd List (2021)

artesunate*	Injection: ampoules, containing 60 mg anhydrous artesunic acid with a separate ampoule of 5% sodium bicarbonate solution. For use in the management of severe malaria. Rectal dosage form: 50 mg [c]; 100 mg [c]; 200 mg capsules (for pre-referral treatment of severe malaria only; patients should be taken to an appropriate health facility for follow-up care) [c]. Tablet: 50 mg. *To be used in combination with either amodiaquine, mefloquine or sulfadoxine + pyrimethamine.
artesunate + amodiaquine*	Tablet: 25 mg + 67.5 mg; 50 mg + 135 mg; 100 mg + 270 mg. *Other combinations that deliver the target doses required such as 153 mg or 200 mg (as hydrochloride) with 50 mg artesunate can be alternatives.
artesunate + mefloquine	Tablet: 25 mg + 55 mg; 100 mg + 220 mg.
artesunate + pyronaridine tetraphosphate ^[a]	Granules: 20 mg + 60 mg [c]. Tablet: 60 mg + 180 mg. ^[a] > 5 kg
chloroquine*	Oral liquid: 50 mg/5 mL (as phosphate or sulfate). Tablet: 100 mg; 150 mg (as phosphate or sulfate). *For use only for the treatment of <i>Plasmodium vivax</i> infection.
dihydroartemisinin + piperaquine phosphate ^[a]	Tablet: 20 mg + 160 mg; 40 mg + 320 mg. ^[a] > 5 kg
doxycycline*	Capsule: 100 mg (as hydrochloride or hyclate). Tablet (dispersible): 100 mg (as monohydrate). *For use only in combination with quinine.
mefloquine*	Tablet: 250 mg (as hydrochloride). *To be used in combination with artesunate 50 mg.
primaquine*	Tablet: 7.5 mg; 15 mg (as diphosphate). *Only for use to achieve radical cure of <i>Plasmodium vivax</i> and <i>Plasmodium ovale</i> infections, given for 14 days.
quinine*	Injection: 300 mg/mL (hydrochloride) in 2 mL ampoule. Tablet: 300 mg (sulfate) or 300 mg (bisulfate). *For use only in the management of severe malaria and should be used in combination with doxycycline.
sulfadoxine + pyrimethamine*	Tablet: 500 mg + 25 mg. *Only in combination with artesunate 50 mg.

WHO Model List of Essential Medicines – 22nd List (2021)

6.5.3.2 For chemoprevention	
amodiaquine – sulfadoxine + pyrimethamine [c]	Co-packaged dispersible tablets: amodiaquine 76.5 mg (as hydrochloride) [3] and sulfadoxine + pyrimethamine 250 mg + 12.5 mg [1]; amodiaquine 153 mg (as hydrochloride) [3] and sulfadoxine + pyrimethamine 500 mg + 25 mg [1].
chloroquine*	Oral liquid: 50 mg/5 mL (as phosphate or sulfate). Tablet: 150 mg (as phosphate or sulfate). *For use only in central American regions, for <i>Plasmodium vivax</i> infections.
doxycycline[a]	Solid oral dosage form: 100 mg (as hydrochloride or hyclate). [a] > 8 years.
mefloquine[a]	Tablet: 250 mg (as hydrochloride). [a] > 5 kg or > 3 months.
proguanil*	Tablet: 100 mg (as hydrochloride). *For use only in combination with chloroquine.
sulfadoxine + pyrimethamine	Tablet: 250 mg + 12.5 mg [c]; 500 mg + 25 mg.
6.5.4 Antipneumocystosis and antitoxoplasmosis medicines	
pyrimethamine	Tablet: 25 mg.
sulfadiazine	Tablet: 500 mg.
sulfamethoxazole + trimethoprim	Injection: 80 mg + 16 mg/mL in 5 mL ampoule; 80 mg + 16 mg/mL in 10 mL ampoule. Oral liquid: 200 mg + 40 mg/5 mL [c]. Tablet: 100 mg + 20 mg; 400 mg + 80 mg [c]; 800 mg + 160 mg
Complementary List	
pentamidine	Tablet: 200 mg; 300 mg (as isethionate).
6.5.5 Antitrypanosomal medicines	
6.5.5.1 African trypanosomiasis	
fexinidazole*	Tablet: 600 mg *For the treatment of 1 st and 2 nd stage of human African trypanosomiasis due to <i>Trypanosoma brucei gambiense</i> infection.
Medicines for the treatment of 1 st stage African trypanosomiasis	
pentamidine*	Powder for injection: 200 mg (as isethionate) in vial. *To be used for the treatment of <i>Trypanosoma brucei gambiense</i> infection.
suramin sodium*	Powder for injection: 1 g in vial. *To be used for the treatment of the initial phase of <i>Trypanosoma brucei rhodesiense</i> infection.

WHO Model List of Essential Medicines – 22nd List (2021)

Medicines for the treatment of 2 nd stage African trypanosomiasis	
eflornithine*	Injection: 200 mg/mL (hydrochloride) in 100 mL bottle. *To be used for the treatment of <i>Trypanosoma brucei gambiense</i> infection.
melarsoprol	Injection: 180 mg/5 mL in 5 mL ampoule (3.6% solution).
nifurtimox *	Tablet: 120 mg. *Only to be used in combination with eflornithine, for the treatment of <i>Trypanosoma brucei gambiense</i> infection.
Complementary List	
melarsoprol [c]	Injection: 180 mg/5 mL in 5 mL ampoule (3.6% solution).
6.5.5.2 American trypanosomiasis	
benznidazole	Tablet: 12.5 mg [c]; 100 mg. Tablet (scored): 50 mg.
nifurtimox	Tablet: 30 mg; 120 mg; 250 mg.
6.6 Medicines for ectoparasitic infections	
ivermectin	Tablet (scored): 3 mg
7. ANTIMIGRAINE MEDICINES	
7.1 For treatment of acute attack	
acetylsalicylic acid	Tablet: 300 mg to 500 mg.
ibuprofen [c]	Tablet: 200 mg; 400 mg.
paracetamol	Oral liquid: 120 mg/5 mL [c]; 125 mg/5 mL [c]. Tablet: 300 mg to 500 mg.
sumatriptan	Tablet: 50 mg
7.2 For prophylaxis	
□ propranolol Therapeutic alternatives to be reviewed (2023)	Tablet: 20 mg; 40 mg (hydrochloride).

WHO Model List of Essential Medicines – 22nd List (2021)

8. IMMUNOMODULATORS AND ANTINEOPLASTICS	
8.1 Immunomodulators for non-malignant disease	
Complementary List	
☐ <i>adalimumab</i> * Therapeutic alternatives*: - certolizumab pegol - etanercept - golimumab - infliximab *including quality-assured biosimilars	Injection: 40 mg/0.8 mL; 40 mg/0.4 mL.
<i>azathioprine</i>	Powder for injection: 100 mg (as sodium salt) in vial. Tablet (scored): 50 mg.
<i>ciclosporin</i>	Capsule: 25 mg. Concentrate for injection: 50 mg/mL in 1 mL ampoule.
<i>tacrolimus</i>	Capsule (immediate-release): 0.5 mg; 0.75 mg; 1 mg; 2 mg; 5 mg. Granules for oral suspension: 0.2 mg; 1 mg. Injection: 5 mg/mL in 1 mL vial.
8.2 Antineoplastics and supportive medicines	
Medicines listed below should be used according to protocols for treatment of the diseases.	
8.2.1 Cytotoxic medicines	
Complementary List	
<i>arsenic trioxide</i>	Concentrate for solution for infusion: 1 mg/mL - Acute promyelocytic leukaemia
<i>asparaginase</i> * *including quality-assured biosimilars	Powder for injection: 10 000 IU in vial. - Acute lymphoblastic leukaemia.
<i>bendamustine</i>	Injection: 45 mg/0.5 mL; 180 mg/2 mL. - Chronic lymphocytic leukaemia - Follicular lymphoma
<i>bleomycin</i>	Powder for injection: 15 mg (as sulfate) in vial. - Hodgkin lymphoma - Kaposi sarcoma - Ovarian germ cell tumour - Testicular germ cell tumour
<i>calcium folinate</i>	Injection: 3 mg/mL in 10 mL ampoule. Tablet: 5 mg; 15 mg; 25 mg. - Burkitt lymphoma - Early stage colon cancer - Early stage rectal cancer - Gestational trophoblastic neoplasia - Metastatic colorectal cancer - Osteosarcoma

WHO Model List of Essential Medicines – 22nd List (2021)

capecitabine	Tablet: 150 mg; 500 mg. – Early stage colon cancer – Early stage rectal cancer – Metastatic breast cancer – Metastatic colorectal cancer
carboplatin	Injection: 50 mg/5 mL; 150 mg/15 mL; 450 mg/45 mL; 600 mg/60 mL. – Cervical cancer – Early stage breast cancer – Epithelial ovarian cancer – Head and neck cancer (as a radio-sensitizer) – Low-grade glioma – Nasopharyngeal cancer – Nephroblastoma (Wilms tumour) – Non-small cell lung cancer – Osteosarcoma – Ovarian germ cell tumour – Retinoblastoma – Testicular germ cell tumour
chlorambucil	Tablet: 2 mg. – Chronic lymphocytic leukaemia
cisplatin	Injection: 10 mg/10 mL; 20 mg/20 mL; 50 mg/50 mL; 100 mg/100 mL. – Cervical cancer – Head and neck cancer (as a radio-sensitizer) – Low-grade glioma – Nasopharyngeal cancer (as a radio-sensitizer) – Non-small cell lung cancer – Osteosarcoma – Ovarian germ cell tumour – Testicular germ cell tumour
cyclophosphamide	Powder for injection: 500 mg; 1 g; 2 g in vial. Tablet: 25 mg, 50 mg. – Acute lymphoblastic leukaemia – Burkitt lymphoma – Chronic lymphocytic leukaemia – Diffuse large B-cell lymphoma – Early stage breast cancer – Ewing sarcoma – Follicular lymphoma – Gestational trophoblastic neoplasia – Hodgkin lymphoma – Low-grade glioma – Metastatic breast cancer – Multiple myeloma – Nephroblastoma (Wilms tumour) – Rhabdomyosarcoma
cytarabine	Powder for injection: 100 mg in vial. – Acute lymphoblastic leukaemia – Acute myeloid leukaemia – Acute promyelocytic leukaemia – Burkitt lymphoma

WHO Model List of Essential Medicines – 22nd List (2021)

dacarbazine	Powder for injection: 100 mg in vial. – Hodgkin lymphoma
dactinomycin	Powder for injection: 500 micrograms in vial. – Ewing sarcoma – Gestational trophoblastic neoplasia – Nephroblastoma (Wilms tumour) – Rhabdomyosarcoma
daunorubicin	Powder for injection: 50 mg (hydrochloride) in vial. – Acute lymphoblastic leukaemia – Acute myeloid leukaemia – Acute promyelocytic leukaemia
docetaxel	Injection: 20 mg/mL; 40 mg/mL. – Early stage breast cancer – Metastatic breast cancer – Metastatic prostate cancer
doxorubicin	Powder for injection: 10 mg; 50 mg (hydrochloride) in vial. – Acute lymphoblastic leukaemia – Burkitt lymphoma – Diffuse large B-cell lymphoma – Early stage breast cancer – Ewing sarcoma – Follicular lymphoma – Hodgkin lymphoma – Kaposi sarcoma – Metastatic breast cancer – Multiple myeloma – Nephroblastoma (Wilms tumour) – Osteosarcoma
etoposide	Capsule: 50 mg, 100 mg. Injection: 20 mg/mL in 5 mL ampoule. – Acute lymphoblastic leukaemia – Acute myeloid leukaemia – Burkitt lymphoma – Ewing sarcoma – Gestational trophoblastic neoplasia – Hodgkin lymphoma – Nephroblastoma (Wilms tumour) – Non-small cell lung cancer – Osteosarcoma – Ovarian germ cell tumour – Retinoblastoma – Testicular germ cell tumour
fludarabine	Powder for injection: 50 mg (phosphate) in vial. Tablet: 10 mg – Chronic lymphocytic leukaemia,

WHO Model List of Essential Medicines – 22nd List (2021)

<i>fluorouracil</i>	Injection: 50 mg/mL in 5 mL ampoule. – Early stage breast cancer – Early stage colon cancer – Early stage rectal cancer – Metastatic colorectal cancer – Nasopharyngeal cancer
<i>gemcitabine</i>	Powder for injection: 200 mg; 1 g in vial. – Epithelial ovarian cancer – Non-small cell lung cancer
<i>hydroxycarbamide</i>	Solid oral dosage form: 200 mg; 250 mg; 300 mg; 400 mg; 500 mg; 1 g. – Chronic myeloid leukaemia
<i>ifosfamide</i>	Powder for injection: 500 mg; 1 g; 2 g in vial. – Burkitt lymphoma – Ewing sarcoma – Nephroblastoma (Wilms tumour) – Ovarian germ cell tumour – Osteosarcoma – Rhabdomyosarcoma – Testicular germ cell tumour
<i>irinotecan</i>	Injection: 40 mg/2 mL in 2 mL vial; 100 mg/5 mL in 5 mL vial; 500 mg/25 mL in 25 mL vial. – Metastatic colorectal cancer – Nephroblastoma (Wilms tumour) – Rhabdomyosarcoma
<i>melfalan</i>	Tablet: 2 mg Powder for injection: 50 mg in vial – Multiple myeloma
<i>mercaptopurine</i>	Tablet: 50 mg. – Acute lymphoblastic leukaemia – Acute promyelocytic leukaemia
<i>methotrexate</i>	Powder for injection: 50 mg (as sodium salt) in vial. Tablet: 2.5 mg (as sodium salt). – Acute lymphoblastic leukaemia – Acute promyelocytic leukaemia – Burkitt lymphoma – Early stage breast cancer – Gestational trophoblastic neoplasia – Osteosarcoma
<i>oxaliplatin</i>	Injection: 50 mg/10 mL in 10 mL vial; 100 mg/20 mL in 20 mL vial; 200 mg/40 mL in 40 mL vial. Powder for injection: 50 mg; 100 mg in vial. – Early stage colon cancer – Metastatic colorectal cancer

WHO Model List of Essential Medicines – 22nd List (2021)

<i>paclitaxel</i>	Injection: 6 mg/mL in vial. – Cervical cancer – Epithelial ovarian cancer – Early stage breast cancer – Metastatic breast cancer – Kaposi sarcoma – Nasopharyngeal cancer – Non-small cell lung cancer – Ovarian germ cell tumour
<i>pegaspargase*</i> *including quality-assured biosimilars	Injection: 3,750 units/5 mL in vial. – Acute lymphoblastic leukaemia
<i>procarbazine [c]</i>	Capsule: 50 mg (as hydrochloride). – Hodgkin lymphoma
<i>reiaiger-Indigo naturalis formulation</i>	Tablet: 270 mg (containing tetra-arsenic tetra-sulfide 30 mg). – Acute promyelocytic leukaemia
<i>tioguanine [c]</i>	Solid oral dosage form: 40 mg. – Acute lymphoblastic leukaemia
<i>vinblastine</i>	Injection: 10 mg/10 mL (sulfate) in vial. Powder for injection: 10 mg (sulfate) in vial. – Hodgkin lymphoma – Kaposi sarcoma – Low-grade glioma – Ovarian germ cell tumour – Testicular germ cell tumour
<i>vincristine</i>	Injection: 1 mg/mL (sulfate); 2 mg/2 mL (sulfate) in vial. Powder for injection: 1 mg; 5 mg (sulfate) in vial. – Acute lymphoblastic leukaemia – Burkitt lymphoma – Diffuse large B-cell lymphoma – Ewing sarcoma – Follicular lymphoma – Gestational trophoblastic neoplasia – Hodgkin lymphoma – Kaposi sarcoma – Low-grade glioma – Nephroblastoma (Wilms tumour) – Retinoblastoma – Rhabdomyosarcoma
<i>vinorelbine</i>	Capsule: 20 mg; 30 mg; 80 mg. Injection: 10 mg/mL in 1 mL vial; 50 mg/5 mL in 5 mL vial. – Non-small cell lung cancer – Metastatic breast cancer – Rhabdomyosarcoma

WHO Model List of Essential Medicines – 22nd List (2021)

8.2.2 Targeted therapies	
Complementary List	
all-trans retinoid acid (ATRA)	Capsule: 10 mg. – Acute promyelocytic leukaemia
bortezomib	Powder for injection: 3.5 mg in vial. – Multiple myeloma
dasatinib	Tablet: 20 mg; 50 mg; 70 mg; 80 mg; 100 mg; 140 mg. – Imatinib-resistant chronic myeloid leukaemia
□ erlotinib Therapeutic alternatives: – afatinib – gefitinib	Tablet: 100 mg; 150 mg. – EGFR mutation-positive advanced non-small cell lung cancer
everolimus	Tablet: 2.5 mg; 5 mg; 7.5 mg; 10 mg. Tablet (dispersible): 2 mg; 3 mg; 5 mg. – Subependymal giant cell astrocytoma
ibrutinib	Capsule: 140 mg. – Relapsed/refractory chronic lymphocytic leukaemia
imatinib	Solid oral dosage form: 100 mg; 400 mg. – Chronic myeloid leukaemia – Gastrointestinal stromal tumour – Philadelphia chromosome positive acute lymphoblastic leukaemia
nilotinib	Capsule: 150 mg; 200 mg. – Imatinib-resistant chronic myeloid leukaemia
rituximab* *including quality-assured biosimilars	Injection (intravenous): 100 mg/10 mL in 10 mL vial; 500 mg/50 mL in 50 mL vial. – Diffuse large B-cell lymphoma – Chronic lymphocytic leukaemia – Follicular lymphoma
trastuzumab* *including quality-assured biosimilars	Powder for injection: 60 mg; 150 mg; 440 mg in vial. – Early stage HER2 positive breast cancer – Metastatic HER2 positive breast cancer

WHO Model List of Essential Medicines – 22nd List (2021)

8.2.3 Immunomodulators	
Complementary List	
<i>filgrastim*</i> <i>*including quality-assured biosimilars</i>	Injection: 120 micrograms/0.2 mL; 300 micrograms/0.5 mL; 480 micrograms/0.8 mL in pre-filled syringe. Injection: 300 micrograms/mL in 1 mL vial; 480 micrograms/1.6 mL in 1.6 mL vial. – Primary prophylaxis in patients at high risk for developing febrile neutropenia associated with myelotoxic chemotherapy. – Secondary prophylaxis for patients who have experienced neutropenia following prior myelotoxic chemotherapy. – To facilitate administration of dose dense chemotherapy regimens.
<i>lenalidomide</i>	Capsule: 25 mg. – Multiple myeloma
☐ <i>nivolumab*</i> Therapeutic alternatives*: – pembrolizumab <i>*including quality-assured biosimilars</i>	Concentrate solution for infusion: 10 mg/mL. – Metastatic melanoma
<i>thalidomide</i>	Capsule: 50 mg. – Multiple myeloma
8.2.4 Hormones and antihormones	
Complementary List	
☐ <i>abiraterone</i> Therapeutic alternatives: – enzalutamide	Tablet: 250 mg; 500 mg. – Metastatic castration-resistant prostate cancer
☐ <i>anastrozole</i> Therapeutic alternatives: – 4 th level ATC chemical subgroup (L02BG Aromatase inhibitors)	Tablet: 1 mg. – Early stage breast cancer – Metastatic breast cancer
☐ <i>bicalutamide</i> Therapeutic alternatives: – flutamide – nilutamide	Tablet: 50 mg. – Metastatic prostate cancer
<i>dexamethasone</i>	Injection: 4 mg/mL (as disodium phosphate salt) in 1 mL ampoule. Oral liquid: 2 mg/5 mL [c]. Tablet: 2 mg [c]; 4 mg. – Acute lymphoblastic leukaemia – Burkitt lymphoma – Multiple myeloma
<i>hydrocortisone</i>	Powder for injection: 100 mg (as sodium succinate) in vial. – Acute lymphoblastic leukaemia – Burkitt lymphoma

WHO Model List of Essential Medicines – 22nd List (2021)

☐ leuprorelin Therapeutic alternatives: - goserelin - triptorelin	Injection: 7.5 mg; 22.5 mg in pre-filled syringe. - Early stage breast cancer - Metastatic prostate cancer.
methylprednisolone [c]	Injection: 40 mg/mL (as sodium succinate) in 1 mL single-dose vial and 5 mL multi-dose vials; 80 mg/mL (as sodium succinate) in 1 mL single-dose vial. - Acute lymphoblastic leukaemia - Burkitt lymphoma
☐ prednisolone Therapeutic alternatives: - prednisone	Oral liquid: 5 mg/mL [c]. Tablet: 5 mg; 25 mg. - Acute lymphoblastic leukaemia - Burkitt lymphoma - Chronic lymphocytic leukaemia - Diffuse large B-cell lymphoma - Follicular lymphoma - Hodgkin lymphoma - Metastatic castration-resistant prostate cancer - Multiple myeloma
tamoxifen	Tablet: 10 mg; 20 mg (as citrate). - Early stage breast cancer - Metastatic breast cancer.
8.2.5 Supportive medicines	
Complementary List	
allopurinol [c]	Tablet: 100 mg; 300 mg. - Tumour lysis syndrome
mesna	Injection: 100 mg/mL in 4 mL and 10 mL ampoules. Tablet: 400 mg; 600 mg. - Burkitt lymphoma - Ewing sarcoma - Nephroblastoma (Wilms tumour) - Ovarian germ cell tumour - Osteosarcoma - Rhabdomyosarcoma - Testicular germ cell tumour
rasburicase	Powder and solvent for solution for infusion: 1.5 mg; 7.5 mg in vial - Tumour lysis syndrome
zoledronic acid	Concentrate solution for infusion: 4 mg/5 mL in 5 mL vial. Solution for infusion: 4 mg/100 mL in 100 mL bottle. - Malignancy-related bone disease

WHO Model List of Essential Medicines – 22nd List (2021)

9. ANTIPARKINSONISM MEDICINES	
☐ biperiden Therapeutic alternatives: – trihexyphenidyl	Injection: 5 mg (lactate) in 1 mL ampoule. Tablet: 2 mg (hydrochloride).
levodopa + ☐ carbidopa Therapeutic alternatives: – benserazide (for carbidopa)	Tablet: 100 mg + 10 mg; 100 mg + 25 mg; 250 mg + 25 mg.
10. MEDICINES AFFECTING THE BLOOD	
10.1 Antianaemia medicines	
ferrous salt	Oral liquid: equivalent to 25 mg iron (as sulfate)/mL. Tablet: equivalent to 60 mg iron.
ferrous salt + folic acid	Tablet: equivalent to 60 mg iron + 400 micrograms folic acid. *nutritional supplement for use during pregnancy
folic acid	Tablet: 400 micrograms*; 1 mg; 5 mg. *periconceptual use for prevention of first occurrence of neural tube defects
hydroxocobalamin	Injection: 1 mg/mL (as acetate, as hydrochloride or as sulfate) in 1 mL ampoule.
<i>Complementary List</i>	
☐ erythropoiesis-stimulating agents* Therapeutic alternatives: - epoetin alfa, beta and theta - darbepoetin alfa - methoxy polyethylene glycol-epoetin beta *including quality-assured biosimilars	Injection: pre-filled syringe 1000 IU/0.5 mL; 2000 IU/0.5 mL; 3000 IU/0.3 mL; 4000 IU/0.4 mL; 5000 IU/0.5 mL; 6000 IU/0.6 mL; 8000 IU/0.8 mL; 10 000 IU/1 mL; 20 000 IU/0.5 mL; 40 000 IU/1 mL.
10.2 Medicines affecting coagulation	
☐ dabigatran Therapeutic alternatives: - apixaban - edoxaban - rivaroxaban	Capsule: 110 mg; 150 mg.
☐ enoxaparin* Therapeutic alternatives*: - dalteparin - nadroparin *including quality-assured biosimilars	Injection: ampoule or pre-filled syringe 20 mg/0.2 mL; 40 mg/0.4 mL; 60 mg/0.6 mL; 80 mg/0.8 mL; 100 mg/1 mL; 120 mg/0.8 mL; 150 mg/1 mL.
heparin sodium	Injection: 1000 IU/mL; 5000 IU/mL; 20 000 IU/mL in 1 mL ampoule.
phytomenadione	Injection: 1 mg/mL [c]; 10 mg/mL in ampoule. Tablet: 10 mg.
protamine sulfate	Injection: 10 mg/mL in 5 mL ampoule.

WHO Model List of Essential Medicines – 22nd List (2021)

tranexamic acid	Injection: 100 mg/mL in 10 mL ampoule.
□ warfarin Therapeutic alternatives to be reviewed (2023)	Tablet: 1 mg; 2 mg; 5 mg (sodium).
<i>Complementary List</i>	
desmopressin[c]	Injection: 4 micrograms/mL (as acetate) in 1 mL ampoule. Nasal spray: 10 micrograms (as acetate) per dose.
heparin sodium [c]	Injection: 1000 IU/mL; 5000 IU/mL in 1 mL ampoule.
protamine sulfate [c]	Injection: 10 mg/mL in 5 mL ampoule.
□ warfarin [c] Therapeutic alternatives to be reviewed (2023)	Tablet: 0.5 mg; 1 mg; 2 mg; 5 mg (sodium).
10.3 Other medicines for haemoglobinopathies	
<i>Complementary List</i>	
□ deferoxamine Therapeutic alternatives: - deferasirox (oral)	Powder for injection: 500 mg (mesilate) in vial.
hydroxycarbamide	Solid oral dosage form: 200 mg; 500 mg; 1 g.
11. BLOOD PRODUCTS OF HUMAN ORIGIN AND PLASMA SUBSTITUTES	
11.1 Blood and blood components	
In accordance with the World Health Assembly resolution WHA63.12, WHO recognizes that achieving self-sufficiency, unless special circumstances preclude it, in the supply of safe blood components based on voluntary, non-remunerated blood donation, and the security of that supply are important national goals to prevent blood shortages and meet the transfusion requirements of the patient population. All preparations should comply with the WHO requirements.	
fresh-frozen plasma	
platelets	
red blood cells	
whole blood	
11.2 Plasma-derived medicines	
All human plasma-derived medicines should comply with the WHO requirements.	
11.2.1 Human immunoglobulins	
anti-D immunoglobulin	Injection: 250 micrograms in single-dose vial.
anti-rabies immunoglobulin	Injection: 150 IU/mL in vial.
anti-tetanus immunoglobulin	Injection: 500 IU in vial.

WHO Model List of Essential Medicines – 22nd List (2021)

<i>Complementary List</i>	
<i>normal immunoglobulin</i>	Intramuscular administration: 16% protein solution. * Intravenous administration: 5%; 10% protein solution. ** Subcutaneous administration: 15%; 16% protein solution. * *Indicated for primary immune deficiency. **Indicated for primary immune deficiency and Kawasaki disease.
11.2.2 Blood coagulation factors	
<i>Complementary List</i>	
☐ coagulation factor VIII Therapeutic alternatives to be reviewed (2023)	Powder for injection: 500 IU/vial.
☐ coagulation factor IX Therapeutic alternatives to be reviewed (2023)	Powder for injection: 500 IU/vial, 1000 IU/vial.
11.3 Plasma substitutes	
☐ dextran 70 Therapeutic alternatives: - Polygeline injectable solution 3.5%	Injectable solution: 6%.
12. CARDIOVASCULAR MEDICINES	
12.1 Antianginal medicines	
☐ bisoprolol Therapeutic alternatives: - carvedilol - metoprolol	Tablet: 1.25 mg; 5 mg.
glyceryl trinitrate	Tablet (sublingual): 500 micrograms.
isosorbide dinitrate	Tablet (sublingual): 5 mg.
verapamil	Tablet: 40 mg; 80 mg (hydrochloride).
12.2 Antiarrhythmic medicines	
☐ bisoprolol Therapeutic alternatives: - carvedilol - metoprolol	Tablet: 1.25 mg; 5 mg.
digoxin	Injection: 250 micrograms/mL in 2 mL ampoule. Oral liquid: 50 micrograms/mL. Tablet: 62.5 micrograms; 250 micrograms.
epinephrine (adrenaline)	Injection: 100 micrograms/mL (as acid tartrate or hydrochloride) in 10 mL ampoule.
lidocaine	Injection: 20 mg/mL (hydrochloride) in 5 mL ampoule.
verapamil	Injection: 2.5 mg/mL (hydrochloride) in 2 mL ampoule. Tablet: 40 mg; 80 mg (hydrochloride).

WHO Model List of Essential Medicines – 22nd List (2021)

<i>Complementary List</i>	
amiodarone	Injection: 50 mg/mL (hydrochloride) in 3 mL ampoule. Tablet: 100 mg; 200 mg; 400 mg (hydrochloride).
12.3 Antihypertensive medicines	
☐ amlodipine Therapeutic alternatives: - 4 th level ATC chemical subgroup (C08CA Dihydropyridine derivatives)	Tablet: 5 mg (as maleate, mesylate or besylate).
☐ bisoprolol Therapeutic alternatives: - atenolol* - carvedilol - metoprolol	Tablet: 1.25 mg; 5 mg. *atenolol should not be used as a first-line agent in uncomplicated hypertension in patients > 60 years.
☐ enalapril Therapeutic alternatives: - 4 th level ATC chemical subgroup (C09AA ACE inhibitors, plain)	Tablet: 2.5 mg; 5 mg (as hydrogen maleate).
hydralazine*	Powder for injection: 20 mg (hydrochloride) in ampoule. Tablet: 25 mg; 50 mg (hydrochloride). *Hydralazine is listed for use only in the acute management of severe pregnancy-induced hypertension. Its use in the treatment of essential hypertension is not recommended in view of the evidence of greater efficacy and safety of other medicines.
☐ hydrochlorothiazide Therapeutic alternatives: - chlorothiazide - chlorthalidone - indapamide	Oral liquid: 50 mg/5 mL. Solid oral dosage form: 12.5 mg; 25 mg.
☐ lisinopril + ☐ amlodipine Therapeutic alternatives: - 4 th level ATC chemical subgroup (C09AA ACE inhibitors, plain) (for lisinopril) - 4 th level ATC chemical subgroup (C08CA Dihydropyridine derivatives) (for amlodipine)	Tablet: 10 mg + 5 mg; 20 mg + 5 mg; 20 mg + 10 mg.
☐ lisinopril + ☐ hydrochlorothiazide Therapeutic alternatives: - 4 th level ATC chemical subgroup (C09AA ACE inhibitors, plain) (for lisinopril) - chlorthalidone, chlorothiazide, indapamide (for hydrochlorothiazide)	Tablet: 10 mg + 12.5 mg; 20 mg + 12.5 mg; 20 mg + 25 mg.
☐ losartan Therapeutic alternatives: - 4 th level ATC chemical subgroup (C09CA Angiotensin II receptor blockers (ARBs), plain)	Tablet: 25 mg; 50 mg; 100 mg.

WHO Model List of Essential Medicines – 22nd List (2021)

methyldopa*	Tablet: 250 mg. *Methyldopa is listed for use only in the management of pregnancy-induced hypertension. Its use in the treatment of essential hypertension is not recommended in view of the evidence of greater efficacy and safety of other medicines.
☐ telmisartan + ☐ amlodipine Therapeutic alternatives: - 4 th level ATC chemical subgroup (C09CA Angiotensin II receptor blockers (ARBs), plain) (for telmisartan) - 4 th level ATC chemical subgroup (C08CA Dihydropyridine derivatives) (for amlodipine)	Tablet: 40 mg + 5 mg; 80 mg + 5 mg; 80 mg + 10 mg.
☐ telmisartan + ☐ hydrochlorothiazide Therapeutic alternatives: - 4 th level ATC chemical subgroup (C09CA Angiotensin II receptor blockers (ARBs), plain) (for telmisartan) - chlorthalidone, chlorothiazide, indapamide (for hydrochlorothiazide)	Tablet: 40 mg + 12.5 mg; 80 mg + 12.5 mg; 80 mg + 25 mg.
<i>Complementary List</i>	
sodium nitroprusside	Powder for infusion; 50 mg in ampoule.
12.4 Medicines used in heart failure	
☐ bisoprolol Therapeutic alternatives: - carvedilol - metoprolol	Tablet: 1.25 mg; 5 mg.
digoxin	Injection: 250 micrograms/mL in 2 mL ampoule. Oral liquid: 50 micrograms/mL. Tablet: 62.5 micrograms; 250 micrograms.
☐ enalapril Therapeutic alternatives: - 4 th level ATC chemical subgroup (C09AA ACE inhibitors, plain)	Tablet: 2.5 mg; 5 mg (as hydrogen maleate).
☐ furosemide Therapeutic alternatives: - bumetanide - torasemide	Injection: 10 mg/mL in 2 mL ampoule. Oral liquid: 20 mg/5 mL [c]. Tablet: 40 mg.
☐ hydrochlorothiazide Therapeutic alternatives: - chlorothiazide - chlorthalidone - indapamide	Oral liquid: 50 mg/5 mL. Solid oral dosage form: 25 mg.
☐ losartan Therapeutic alternatives: - 4 th level ATC chemical subgroup (C09CA Angiotensin II receptor blockers (ARBs), plain)	Tablet: 25 mg; 50 mg; 100 mg.

WHO Model List of Essential Medicines – 22nd List (2021)

spironolactone	Tablet: 25 mg.
<i>Complementary List</i>	
dopamine	Injection: 40 mg/mL (hydrochloride) in 5 mL vial.
12.5 Antithrombotic medicines	
12.5.1 Anti-platelet medicines	
acetylsalicylic acid	Tablet: 100 mg.
clopidogrel	Tablet: 75 mg; 300 mg
12.5.2 Thrombolytic medicines	
<i>Complementary List</i>	
alteplase	Powder for injection: 10 mg; 20 mg; 50 mg in vial
streptokinase	Powder for injection: 1.5 million IU in vial.
12.6 Lipid-lowering agents	
☐ simvastatin* Therapeutic alternatives: - atorvastatin - fluvastatin - lovastatin - pravastatin	Tablet: 5 mg; 10 mg; 20 mg; 40 mg. *For use in high-risk patients.
13. DERMATOLOGICAL MEDICINES (topical)	
13.1 Antifungal medicines	
☐ miconazole Therapeutic alternatives: - 4 th level ATC chemical subgroup (D01AC Imidazole and triazole derivatives) excluding combinations	Cream or ointment: 2% (nitrate).
selenium sulfide	Detergent-based suspension: 2%.
sodium thiosulfate	Solution: 15%.
terbinafine	Cream or ointment: 1% (hydrochloride).

WHO Model List of Essential Medicines – 22nd List (2021)

13.2 Anti-infective medicines	
mupirocin	Cream: 2% (as calcium). Ointment: 2%.
potassium permanganate	Aqueous solution: 1:10 000.
silver sulfadiazine ^[a]	Cream: 1%. ^[a] > 2 months.
13.3 Anti-inflammatory and antipruritic medicines	
☐ betamethasone ^[a] Therapeutic alternatives: - 4 th level ATC chemical subgroup (D07AC Corticosteroids, potent (group III))	Cream or ointment: 0.1% (as valerate). ^[a] Hydrocortisone preferred in neonates.
calamine	Lotion.
☐ hydrocortisone Therapeutic alternatives: - 4 th level ATC chemical subgroup (D07AA Corticosteroids, weak (group I))	Cream or ointment: 1% (acetate).
13.4 Medicines affecting skin differentiation and proliferation	
benzoyl peroxide	Cream or lotion: 5%.
☐ calcipotriol Therapeutic alternatives: - calcitriol - tacalcitol	Cream or ointment: 50 micrograms/mL (0.005%). Lotion: 50 micrograms/mL (0.005%).
coal tar	Solution: 5%.
fluorouracil	Ointment: 5%.
☐ podophyllum resin Therapeutic alternatives: - podophylotoxin	Solution: 10% to 25%.
salicylic acid	Solution: 5%.
urea	Cream or ointment: 5%; 10%.
13.5 Scabicides and pediculicides	
☐ benzyl benzoate ^[a] Therapeutic alternatives: - precipitated sulfur topical ointment	Lotion: 25%. ^[a] > 2 years.
permethrin	Cream: 5%. Lotion: 1%.

WHO Model List of Essential Medicines – 22nd List (2021)

14. DIAGNOSTIC AGENTS	
14.1 Ophthalmic medicines	
fluorescein	Eye drops: 1% (sodium salt).
☐ tropicamide Therapeutic alternatives: - atropine - cyclopentolate	Eye drops: 0.5%.
14.2 Radiocontrast media	
☐ amidotrizoate Therapeutic alternatives to be reviewed (2023)	Injection: 140 mg to 420 mg iodine/mL (as sodium or meglumine salt) in 20 mL ampoule.
barium sulfate	Aqueous suspension.
☐ iohexol Therapeutic alternatives to be reviewed (2023)	Injection: 140 mg to 350 mg iodine/mL in 5 mL; 10 mL; 20 mL ampoules.
<i>Complementary List</i>	
barium sulfate [c]	Aqueous suspension.
☐ meglumine iotroxate Therapeutic alternatives to be reviewed (2023)	Solution: 5 g to 8 g iodine in 100 mL to 250 mL.
15. ANTISEPTICS AND DISINFECTANTS	
15.1 Antiseptics	
☐ chlorhexidine Therapeutic alternatives to be reviewed (2023)	Solution: 5% (digluconate).
☐ ethanol Therapeutic alternatives: - propanol	Solution: 70% (denatured).
☐ povidone iodine Therapeutic alternatives: - iodine	Solution: 10% (equivalent to 1% available iodine).
15.2 Disinfectants	
alcohol based hand rub	Solution: containing ethanol 80% volume/volume. Solution: containing isopropyl alcohol 75% volume/volume.
chlorine base compound	Liquid: (0.1% available chlorine) for solution. Powder: (0.1% available chlorine) for solution. Solid: (0.1% available chlorine) for solution.

WHO Model List of Essential Medicines – 22nd List (2021)

☐ chloroxylenol Therapeutic alternatives: - 4 th level ATC chemical subgroup (D08AE Phenol and derivatives)	Solution: 4.8%.
glutaral	Solution: 2%.
16. DIURETICS	
amiloride	Tablet: 5 mg (hydrochloride).
☐ furosemide Therapeutic alternatives: - bumetanide - torasemide	Injection: 10 mg/mL in 2 mL ampoule. Oral liquid: 20 mg/5 mL [c]. Tablet: 10 mg [c]; 20 mg [c]; 40 mg.
☐ hydrochlorothiazide Therapeutic alternatives: - chlorothiazide - chlortalidone - indapamide	Solid oral dosage form: 25 mg.
mannitol	Injectable solution: 10%; 20%.
spironolactone	Tablet: 25 mg.
<i>Complementary List</i>	
☐ hydrochlorothiazide[c] Therapeutic alternatives: - chlorothiazide - chlortalidone	Tablet (scored): 25 mg.
mannitol [c]	Injectable solution: 10%; 20%.
spironolactone[c]	Oral liquid: 5 mg/5 mL; 10 mg/5 mL; 25 mg/5 mL. Tablet: 25 mg.
17. GASTROINTESTINAL MEDICINES	
<i>Complementary List</i>	
pancreatic enzymes[c]	Age-appropriate formulations and doses including lipase, protease and amylase.
17.1 Antiulcer medicines	
☐ omeprazole Therapeutic alternatives: - 4 th level ATC chemical subgroup (A02BG Proton pump inhibitors) excluding combinations	Powder for injection: 40 mg in vial Powder for oral liquid: 20 mg; 40 mg sachets. Solid oral dosage form: 10 mg; 20 mg; 40 mg.
☐ ranitidine Therapeutic alternatives: - 4 th level ATC chemical subgroup (A02BA H ₂ -receptor antagonists) excluding combinations	Injection: 25 mg/mL (as hydrochloride) in 2 mL ampoule. Oral liquid: 75 mg/5 mL (as hydrochloride). Tablet: 150 mg (as hydrochloride).

WHO Model List of Essential Medicines – 22nd List (2021)

17.2 Antiemetic medicines	
dexamethasone	Injection: 4 mg/mL (as disodium phosphate salt) in 1 mL ampoule. Oral liquid: 0.5 mg/5 mL; 2 mg/5 mL. Solid oral dosage form: 0.5 mg; 0.75 mg; 1.5 mg; 4 mg.
metoclopramide ^a	Injection: 5 mg/mL (hydrochloride) in 2 mL ampoule. Oral liquid: 5 mg/5 mL [c]. Tablet: 10 mg (hydrochloride). ^a Not in neonates.
☐ ondansetron ^a Therapeutic alternatives: - dolasetron - granisetron - palonosetron - tropisetron	Injection: 2 mg base/mL in 2 mL ampoule (as hydrochloride). Oral liquid: 4 mg base/5 mL. Solid oral dosage form: Eq 4 mg base; Eq 8 mg base; Eq 24 mg base. ^a > 1 month.
Complementary list	
aprepitant	Capsule: 80 mg; 125 mg; 165 mg Powder for oral suspension: 125 mg in sachet
17.3 Anti-inflammatory medicines	
☐ sulfasalazine Therapeutic alternatives: - mesalazine	Retention enema. Suppository: 500 mg. Tablet: 500 mg.
Complementary List	
hydrocortisone	Retention enema: 100 mg/60 mL. Suppository: 25 mg (acetate).
prednisolone	Retention enema: 20 mg/100 mL (as sodium phosphate).
17.4 Laxatives	
☐ senna Therapeutic alternatives: - bisacodyl	Tablet: 7.5 mg (sennosides) (or traditional dosage forms).
17.5 Medicines used in diarrhoea	
oral rehydration salts – zinc sulfate [c]	Co-package containing: ORS powder for dilution (see Section 17.5.1) – zinc sulfate solid oral dosage form 20 mg (see Section 17.5.2)

WHO Model List of Essential Medicines – 22nd List (2021)

17.5.1 Oral rehydration																					
oral rehydration salts	Powder for dilution in 200 mL; 500 mL; 1 L. <table> <tr> <td>glucose:</td><td>75 mEq</td></tr> <tr> <td>sodium:</td><td>75 mEq or mmol/L</td></tr> <tr> <td>chloride:</td><td>65 mEq or mmol/L</td></tr> <tr> <td>potassium:</td><td>20 mEq or mmol/L</td></tr> <tr> <td>citrate:</td><td>10 mmol/L</td></tr> <tr> <td>osmolality:</td><td>245 mOsm/L</td></tr> <tr> <td>glucose:</td><td>13.5 g/L</td></tr> <tr> <td>sodium chloride:</td><td>2.6 g/L</td></tr> <tr> <td>potassium chloride:</td><td>1.5 g/L</td></tr> <tr> <td>trisodium citrate dihydrate*:</td><td>2.9 g/L</td></tr> </table> *Trisodium citrate dihydrate may be replaced by sodium hydrogen carbonate (sodium bicarbonate) 2.5 g/L. However, as the stability of this latter formulation is very poor under tropical conditions, it is recommended only when manufactured for immediate use.	glucose:	75 mEq	sodium:	75 mEq or mmol/L	chloride:	65 mEq or mmol/L	potassium:	20 mEq or mmol/L	citrate:	10 mmol/L	osmolality:	245 mOsm/L	glucose:	13.5 g/L	sodium chloride:	2.6 g/L	potassium chloride:	1.5 g/L	trisodium citrate dihydrate*:	2.9 g/L
glucose:	75 mEq																				
sodium:	75 mEq or mmol/L																				
chloride:	65 mEq or mmol/L																				
potassium:	20 mEq or mmol/L																				
citrate:	10 mmol/L																				
osmolality:	245 mOsm/L																				
glucose:	13.5 g/L																				
sodium chloride:	2.6 g/L																				
potassium chloride:	1.5 g/L																				
trisodium citrate dihydrate*:	2.9 g/L																				
17.5.2 Medicines for diarrhoea																					
zinc sulfate*	Solid oral dosage form: 20 mg. *In acute diarrhoea zinc sulfate should be used as an adjunct to oral rehydration salts.																				
18. MEDICINES FOR ENDOCRINE DISORDERS																					
18.1 Adrenal hormones and synthetic substitutes																					
fludrocortisone	Tablet: 100 micrograms (acetate).																				
hydrocortisone	Tablet: 5 mg; 10 mg; 20 mg.																				
18.2 Androgens																					
<i>Complementary List</i>																					
testosterone	Injection: 200 mg (enanthate) in 1 mL ampoule.																				
18.3 Estrogens																					
18.4 Progestogens																					
□ medroxyprogesterone acetate Therapeutic alternatives: - norethisterone	Tablet: 5 mg.																				

WHO Model List of Essential Medicines – 22nd List (2021)

18.5 Medicines for diabetes	
18.5.1 Insulins	
insulin injection (soluble)* *including quality-assured biosimilars	Injection: 40 IU/mL in 10 mL vial; 100 IU/mL in 10 mL vial.
intermediate-acting insulin* *including quality-assured biosimilars	Injection: 40 IU/mL in 10 mL vial; 100 IU/mL in 10 mL vial (as compound insulin zinc suspension or isophane insulin).
☐ long-acting insulin analogues* Therapeutic alternatives: - insulin degludec - insulin detemir - insulin glargine *including quality-assured biosimilars	Injection: 100 IU/mL in 3 mL cartridge or pre-filled pen.
18.5.2 Oral hypoglycaemic agents	
☐ empagliflozin Therapeutic alternatives: - canagliflozin - dapagliflozin	Tablet: 10 mg; 25 mg.
☐ gliclazide* Therapeutic alternatives: - 4 th level ATC chemical subgroup (A10BB Sulfonylureas)	Solid oral dosage form: (controlled-release tablets) 30 mg; 60 mg; 80 mg. *glibenclamide not suitable above 60 years.
metformin	Tablet: 500 mg (hydrochloride).
Complementary List	
metformin [c]	Tablet: 500 mg (hydrochloride).
18.6 Medicines for hypoglycaemia	
glucagon	Injection: 1 mg/mL.
Complementary List	
diazoxide [c]	Oral liquid: 50 mg/mL. Tablet: 50 mg.
18.7 Thyroid hormones and antithyroid medicines	
levothyroxine	Tablet: 25 micrograms [c]; 50 micrograms; 100 micrograms (sodium salt).
potassium iodide	Tablet: 60 mg.
☐ methimazole Therapeutic alternatives: - carbimazole (depending on local availability)	Tablet: 5mg, 10mg, 20mg.
propylthiouracil*	Tablet: 50 mg. *For use when alternative first-line treatment is not appropriate or available; and in patients during the first trimester of pregnancy.

WHO Model List of Essential Medicines – 22nd List (2021)

<i>Complementary List</i>	
<i>Lugol's solution [c]</i>	<i>Oral liquid: about 130 mg total iodine/mL.</i>
□ <i>methimazole [c]</i> <i>Therapeutic alternatives:</i> <i>- carbimazole (depending on local availability)</i>	<i>Tablet: 5mg, 10mg, 20mg.</i>
<i>potassium iodide [c]</i>	<i>Tablet: 60 mg.</i>
<i>propylthiouracil* [c]</i>	<i>Tablet: 50 mg.</i> <i>*For use when alternative first-line treatment is not appropriate or available</i>
19. IMMUNOLOGICALS	
19.1 Diagnostic agents	
All tuberculin should comply with the WHO requirements for tuberculin.	
tuberculin, purified protein derivative (PPD)	Injection.
19.2 Sera, immunoglobulins and monoclonal antibodies	
All plasma fractions should comply with the WHO requirements.	
anti-rabies virus monoclonal antibodies* <i>*including quality-assured biosimilars</i>	Injection; 40 IU/mL in 1.25 mL, 2.5 mL vial; 100 IU/mL in 2.5 mL vial (human). Injection; 300 IU/mL in 10 mL vial; 600 IU/mL in 1 mL, 2.5 mL and 5 mL vial (murine).
antivenom immunoglobulin*	Injection. *Exact type to be defined locally.
diphtheria antitoxin	Injection; 10 000 IU; 20 000 IU in vial.
equine rabies immunoglobulin	Injection; 150 IU/mL; 200 IU/mL; 300 IU/mL; 400 IU/mL in vial.

WHO Model List of Essential Medicines – 22nd List (2021)

19.3 Vaccines	
WHO immunization policy recommendations are published in vaccine position papers based on recommendations made by the Strategic Advisory Group of Experts on Immunization (SAGE).	
WHO vaccine position papers are updated three to four times per year. The list below details the vaccines for which there is a recommendation from SAGE and a corresponding WHO position paper as at September 2020. The most recent versions of the WHO position papers, reflecting the current evidence related to a specific vaccine and the related recommendations, can be accessed at any time on the WHO website at: https://www.who.int/teams/immunization-vaccines-and-biologicals/policies/position-papers	
Vaccine recommendations may be universal or conditional (e.g., in certain regions, in some high-risk populations or as part of immunization programmes with certain characteristics). Details are available in the relevant position papers, and in the Summary Tables of WHO Routine Immunization Recommendations available on the WHO website at: https://www.who.int/teams/immunization-vaccines-and-biologicals/policies/who-recommendations-for-routine-immunization---summary-tables	
Selection of vaccines from the Model List will need to be determined by each country after consideration of international recommendations, epidemiology and national priorities.	
All vaccines should comply with the WHO requirements for biological substances.	
WHO noted the need for vaccines used in children to be polyvalent.	
<i>Recommendations for all</i>	
BCG vaccine	
diphtheria vaccine	
Haemophilus influenzae type b vaccine	
hepatitis B vaccine	
human papilloma virus (HPV) vaccine	
measles vaccine	
pertussis vaccine	
pneumococcal vaccine	
poliomyelitis vaccine	
rotavirus vaccine	
rubella vaccine	
tetanus vaccine	
<i>Recommendations for certain regions</i>	
Japanese encephalitis vaccine	
tick-borne encephalitis vaccine	
yellow fever vaccine	
<i>Recommendations for some high-risk populations</i>	
cholera vaccine	
dengue vaccine	
hepatitis A vaccine	
meningococcal meningitis vaccine	

WHO Model List of Essential Medicines – 22nd List (2021)

rabies vaccine	
typhoid vaccine	
<i>Recommendations for immunization programmes with certain characteristics</i>	
influenza vaccine (seasonal)	
mumps vaccine	
varicella vaccine	
20. MUSCLE RELAXANTS (PERIPHERALLY-ACTING) AND CHOLINESTERASE INHIBITORS	
☐ atracurium Therapeutic alternatives to be reviewed (2023)	Injection: 10 mg/mL (besylate).
neostigmine	Injection: 500 micrograms/mL (methylsulfate) in 1 mL ampoule; 2.5 mg/mL (methylsulfate) in 1 mL ampoule. Tablet: 15 mg (bromide).
suxamethonium	Injection: 50 mg/mL (chloride) in 2 mL ampoule. Powder for injection: (chloride), in vial.
☐ vecuronium [c] Therapeutic alternatives to be reviewed (2023)	Powder for injection: 10 mg (bromide) in vial.
<i>Complementary List</i>	
pyridostigmine	Injection: 1 mg in 1 mL ampoule. Tablet: 60 mg (bromide).
☐ vecuronium Therapeutic alternatives to be reviewed (2023)	Powder for injection: 10 mg (bromide) in vial.
21. OPHTHALMOLOGICAL PREPARATIONS	
21.1 Anti-infective agents	
acidovir	Ointment: 3% W/W.
azithromycin	Solution (eye drops): 1.5%. – <i>Trachoma</i>
erythromycin	Ointment: 0.5% [c] – <i>Infections due to Chlamydia trachomatis or Neisseria gonorrhoea.</i>
☐ gentamicin Therapeutic alternatives: – amikacin – kanamycin – netilmicin – tobramycin	Solution (eye drops): 0.3% (sulfate). – <i>Bacterial blepharitis</i> – <i>Bacterial conjunctivitis</i>

WHO Model List of Essential Medicines – 22nd List (2021)

natamycin	Suspension (eye drops): 5% – <i>Fungal keratitis</i>
☐ ofloxacin Therapeutic alternatives: – 4 th level ATC chemical subgroup (S01AE Fluoroquinolones)	Solution (eye drops): 0.3% – <i>Bacterial conjunctivitis</i> – <i>Bacterial keratitis</i>
☐ tetracycline Therapeutic alternatives: – chlortetracycline – oxytetracycline	Eye ointment: 1% (hydrochloride). – <i>Bacterial blepharitis</i> – <i>Bacterial conjunctivitis</i> – <i>Bacterial keratitis</i> – <i>Trachoma</i>
21.2 Anti-inflammatory agents	
☐ prednisolone Therapeutic alternatives to be reviewed (2023)	Solution (eye drops): 0.5% (sodium phosphate).
21.3 Local anaesthetics	
☐ tetracaine ^a Therapeutic alternatives: – 4 th level ATC chemical subgroup (S01HA Local anaesthetics) excluding cocaine and combinations	Solution (eye drops): 0.5% (hydrochloride). ^a Not in preterm neonates.
21.4 Miotics and antiglaucoma medicines	
acetazolamide	Tablet: 250 mg.
latanoprost	Solution (eye drops): 50 micrograms/mL
☐ pilocarpine Therapeutic alternatives: – carbachol	Solution (eye drops): 2%; 4% (hydrochloride or nitrate).
☐ timolol Therapeutic alternatives: – 4 th level ATC chemical subgroup (S01ED Beta blocking agents) excluding combinations	Solution (eye drops): 0.25%; 0.5% (as hydrogen maleate).
21.5 Mydriatics	
☐ atropine ^a Therapeutic alternatives*: – cyclopentolate hydrochloride – homatropine hydrobromide *EMLc only	Solution (eye drops): 0.1%; 0.5%; 1% (sulfate). ^a > 3 months.
Complementary List	
epinephrine (adrenaline)	Solution (eye drops): 2% (as hydrochloride).
21.6 Anti-vascular endothelial growth factor (VEGF) preparations	
Complementary List	
bevacizumab* *including quality-assured biosimilars	Injection: 25 mg/mL

WHO Model List of Essential Medicines – 22nd List (2021)

22. MEDICINES FOR REPRODUCTIVE HEALTH AND PERINATAL CARE	
22.1 Contraceptives	
22.1.1 Oral hormonal contraceptives	
□ ethinylestradiol + □ levonorgestrel Therapeutic alternatives to be reviewed (2023)	Tablet: 30 micrograms + 150 micrograms.
□ ethinylestradiol + □ norethisterone Therapeutic alternatives to be reviewed (2023)	Tablet: 35 micrograms + 1 mg.
levonorgestrel	Tablet: 30 micrograms; 750 micrograms (pack of two); 1.5 mg.
ulipristal	Tablet: 30 mg (as acetate)
22.1.2 Injectable hormonal contraceptives	
estradiol cypionate + medroxyprogesterone acetate	Injection: 5 mg + 25 mg.
medroxyprogesterone acetate	Injection (intramuscular): 150 mg/mL in 1 mL vial. Injection (subcutaneous): 104 mg/0.65 mL in pre-filled syringe or single-dose injection delivery system.
norethisterone enantate	Oily solution: 200 mg/mL in 1 mL ampoule.
22.1.3 Intrauterine devices	
copper-containing device	
levonorgestrel-releasing intrauterine system	Intrauterine system: with reservoir containing 52 mg of levonorelrel
22.1.4 Barrier methods	
condoms	
diaphragms	
22.1.5 Implantable contraceptives	
etonogestrel-releasing implant	Single-rod etonogestrel-releasing implant: containing 68 mg of etonogestrel.
levonorgestrel-releasing implant	Two-rod levonorgestrel-releasing implant: each rod containing 75 mg of levonorgestrel (150 mg total).
22.1.6 Intravaginal contraceptives	
ethinylestradiol + etonogestrel	Vaginal ring: containing 2.27 mg + 11.7 mg
progesterone vaginal ring*	Progesterone-releasing vaginal ring: containing 2.074 g of micronized progesterone. *For use in women actively breastfeeding at least 4 times per day
22.2 Ovulation inducers	
Complementary List	
clomifene	Tablet: 50 mg (citrate).

WHO Model List of Essential Medicines – 22nd List (2021)

22.3 Uterotonics	
carbetocin	Injection (heat stable): 100 micrograms/mL
☐ ergometrine Therapeutic alternatives: - methylergometrine	Injection: 200 micrograms (hydrogen maleate) in 1 mL ampoule.
mifepristone – misoprostol Where permitted under national law and where culturally acceptable.	Tablet 200 mg – tablet 200 micrograms. Co-package containing: mifepristone 200 mg tablet [1] and misoprostol 200 micrograms tablet [4]
misoprostol	Tablet: 200 micrograms. – Management of incomplete abortion and miscarriage; – Prevention and treatment of postpartum haemorrhage where oxytocin is not available or cannot be safely used Vaginal tablet: 25 micrograms.* *Only for use for induction of labour where appropriate facilities are available.
oxytocin	Injection: 10 IU in 1 mL.
22.4 Antioxytotics (tocolytics)	
nifedipine	Immediate-release capsule: 10 mg.
22.5 Other medicines administered to the mother	
dexamethasone	Injection: 4 mg/mL (as disodium phosphate salt) in 1 mL ampoule.
multiple micronutrient supplement	Tablet containing: Vitamin A (retinol acetate) 800 micrograms retinol activity equivalent Vitamin C (ascorbic acid) 70 mg Vitamin D (cholecalciferol) 5 micrograms (200 IU) Vitamin E (alpha tocopherol succinate) 10 mg alpha tocopherol equivalent Vitamin B1 (thiamine mononitrate) 1.4 mg Vitamin B2 (riboflavin) 1.4 mg Vitamin B3 (niacinamide) 18 mg niacin equivalent Vitamin B6 (pyridoxine hydrochloride) 1.9 mg Folic acid (folic acid) 660 micrograms dietary folate equivalent (400 micrograms) Vitamin B12 (cyanocobalamin) 2.6 micrograms Iron (ferrous fumarate) 30 mg Iodine (potassium iodide) 150 micrograms Zinc (zinc oxide) 15 mg Selenium (sodium selenite) 65 micrograms Copper (cupric oxide) 2 mg
tranexamic acid	Injection: 100 mg/mL in 10 mL ampoule
22.6 Medicines administered to the neonate [c]	
caffeine citrate [c]	Injection: 20 mg/mL (equivalent to 10 mg caffeine base/mL). Oral liquid: 20 mg/mL (equivalent to 10 mg caffeine base/mL).
chlorhexidine [c]	Solution or gel: 7.1% (digluconate) delivering 4% chlorhexidine (for umbilical cord care).

WHO Model List of Essential Medicines – 22nd List (2021)

<i>Complementary List</i>	
☐ <i>ibuprofen [c]</i> Therapeutic alternatives: - <i>indometacin</i>	Solution for injection: 5 mg/mL.
☐ <i>prostaglandin E1 [c]</i> Therapeutic alternatives: - <i>prostaglandin E2</i>	Solution for injection: 0.5 mg/mL in alcohol.
<i>surfactant [c]</i>	Suspension for intratracheal instillation: 25 mg/mL or 80 mg/mL.
23. PERITONEAL DIALYSIS SOLUTION	
<i>Complementary List</i>	
<i>intrapertitoneal dialysis solution (of appropriate composition)</i>	<i>Parenteral solution.</i>
24. MEDICINES FOR MENTAL AND BEHAVIOURAL DISORDERS	
24.1 Medicines used in psychotic disorders	
☐ <i>chlorpromazine</i> Therapeutic alternatives to be reviewed (2023)	Injection: 25 mg/mL (hydrochloride) in 2 mL ampoule. Oral liquid: 25 mg/5 mL (hydrochloride). Tablet: 100 mg (hydrochloride).
☐ <i>fluphenazine</i> Therapeutic alternatives to be reviewed (2023)	Injection: 25 mg (decanoate or enantate) in 1 mL ampoule.
☐ <i>haloperidol</i> Therapeutic alternatives to be reviewed (2023)	Injection: 5 mg in 1 mL ampoule. Tablet: 2 mg; 5 mg.
☐ <i>paliperidone</i> Therapeutic alternatives: - <i>risperidone injection</i>	Injection (prolonged-release): 25 mg; 50 mg; 75 mg; 100 mg; 150 mg (as palmitate) in pre-filled syringe
<i>risperidone</i>	Solid oral dosage form: 0.25 mg to 6.0 mg.
<i>Complementary List</i>	
<i>chlorpromazine [c]</i>	Injection: 25 mg/mL (hydrochloride) in 2 mL ampoule. Oral liquid: 25 mg/5 mL (hydrochloride). Tablet: 10 mg; 25 mg; 50 mg; 100 mg (hydrochloride).
<i>clozapine</i>	Solid oral dosage form: 25 to 200 mg.
<i>haloperidol [c]</i>	Injection: 5 mg in 1 mL ampoule. Oral liquid: 2 mg/mL. Solid oral dosage form: 0.5 mg; 2 mg; 5 mg.

WHO Model List of Essential Medicines – 22nd List (2021)

24.2 Medicines used in mood disorders	
24.2.1 Medicines used in depressive disorders	
☐ amitriptyline Therapeutic alternatives to be reviewed (2023)	Tablet: 25 mg; 75mg, (hydrochloride).
☐ fluoxetine Therapeutic alternatives: - citalopram - escitalopram - fluvoxamine - paroxetine - sertraline	Solid oral dosage form: 20 mg (as hydrochloride).
<i>Complementary List</i>	
fluoxetine  [c]	Solid oral dosage form: 20 mg (as hydrochloride).  > 8 years.
24.2.2 Medicines used in bipolar disorders	
carbamazepine	Tablet (scored): 100 mg; 200 mg.
lithium carbonate	Solid oral dosage form: 300 mg.
valproic acid (sodium valproate)* *avoid use in pregnancy and in women and girls of child-bearing potential, unless alternative treatments are ineffective or not tolerated because of the high risk of birth defects and developmental disorders in children exposed to valproate in the womb.	Tablet (enteric-coated): 200 mg; 500 mg.
24.3 Medicines for anxiety disorders	
☐ diazepam Therapeutic alternatives to be reviewed (2023)	Tablet (scored): 2 mg; 5 mg.
24.4 Medicines used for obsessive compulsive disorders	
clomipramine	Capsule: 10 mg; 25 mg (hydrochloride).
24.5 Medicines for disorders due to psychoactive substance use	
bupropion	Tablet (sustained-release): 150 mg (hydrochloride)
nicotine replacement therapy (NRT)	Chewing gum: 2 mg; 4 mg (as polacrilex). Transdermal patch: 5 mg to 30 mg/16 hrs; 7 mg to 21 mg/24 hrs.
varenicline	Tablet: 0.5 mg, 1 mg
<i>Complementary List</i>	
☐ methadone* Therapeutic alternatives: - buprenorphine	Concentrate for oral liquid: 5 mg/mL; 10 mg/mL (hydrochloride). Oral liquid: 5 mg/5 mL; 10 mg/5 mL (hydrochloride). *The medicines should only be used within an established support programme.

WHO Model List of Essential Medicines – 22nd List (2021)

25. MEDICINES ACTING ON THE RESPIRATORY TRACT	
25.1 Antiasthmatic medicines and medicines for chronic obstructive pulmonary disease	
☐ budesonide Therapeutic alternatives: - beclomethasone - ciclesonide - flunisolide - fluticasone - mometasone	Inhalation (aerosol): 100 micrograms per dose; 200 micrograms per dose.
☐ budesonide + ☐ formoterol Therapeutic alternatives: - beclomethasone + formoterol - budesonide + salmeterol - fluticasone + formoterol - fluticasone furoate + vilanterol - mometasone + formoterol	Dry powder inhaler: 100 micrograms + 6 micrograms per dose; 200 micrograms + 6 micrograms per dose
epinephrine (adrenaline)	Injection: 1 mg/mL (as hydrochloride or hydrogen tartrate) in 1 mL ampoule.
ipratropium bromide	Inhalation (aerosol): 20 micrograms/metered dose.
☐ salbutamol Therapeutic alternatives: - terbutaline	Inhalation (aerosol): 100 micrograms (as sulfate) per dose. Injection: 50 micrograms/mL (as sulfate) in 5 mL ampoule. Metered dose inhaler (aerosol): 100 micrograms (as sulfate) per dose. Respirator solution for use in nebulizers: 5 mg/mL (as sulfate).
☐ tiotropium Therapeutic alternatives: - acclidinium - glycopyrronium - umeclidinium	Powder for inhalation, capsule: 18 micrograms Inhalation solution: 1.25 micrograms; 2.5 micrograms per actuation
26. SOLUTIONS CORRECTING WATER, ELECTROLYTE AND ACID-BASE DISTURBANCES	
26.1 Oral	
oral rehydration salts	See section 17.5.1.
potassium chloride	Powder for solution.
26.2 Parenteral	
glucose	Injectable solution: 5% (isotonic); 10% (hypertonic); 50% (hypertonic).
glucose with sodium chloride	Injectable solution: 4% glucose, 0.18% sodium chloride (equivalent to Na ⁺ 30 mmol/L, Cl ⁻ 30 mmol/L). Injectable solution: 5% glucose, 0.9% sodium chloride (equivalent to Na ⁺ 150 mmol/L and Cl ⁻ 150 mmol/L); 5% glucose, 0.45% sodium chloride (equivalent to Na ⁺ 75 mmol/L and Cl ⁻ 75 mmol/L) [c].

WHO Model List of Essential Medicines – 22nd List (2021)

potassium chloride	Solution: 11.2% in 20 mL ampoule (equivalent to K ⁺ 1.5 mmol/mL, Cl ⁻ 1.5 mmol/mL). Solution for dilution: 7.5% (equivalent to K 1 mmol/mL and Cl 1 mmol/mL) [c]; 15% (equivalent to K 2 mmol/mL and Cl 2 mmol/mL) [c].
sodium chloride	Injectable solution: 0.9% isotonic (equivalent to Na ⁺ 154 mmol/L, Cl ⁻ 154 mmol/L).
sodium hydrogen carbonate	Injectable solution: 1.4% isotonic (equivalent to Na ⁺ 167 mmol/L, HCO ₃ ⁻ 167 mmol/L). Solution: 8.4% in 10 mL ampoule (equivalent to Na ⁺ 1000 mmol/L, HCO ₃ ⁻ 1000 mmol/L).
sodium lactate, compound solution	Injectable solution.
26.3 Miscellaneous	
water for injection	2 mL; 5 mL; 10 mL ampoules.
27. VITAMINS AND MINERALS	
ascorbic acid	Tablet: 50 mg.
calcium	Tablet: 500 mg (elemental).
□ colecalciferol [c] Therapeutic alternatives: - ergocalciferol	Oral liquid: 400 IU/mL. Solid oral dosage form: 400 IU, 1000 IU.
□ ergocalciferol Therapeutic alternatives: - colecalciferol	Oral liquid: 250 micrograms/mL (10 000 IU/mL). Solid oral dosage form: 1.25 mg (50 000 IU).
iodine	Capsule: 190 mg. Iodized oil: 1 mL (480 mg iodine); 0.5 mL (240 mg iodine) in ampoule (oral or injectable); 0.57 mL (308 mg iodine) in dispenser bottle.
multiple micronutrient powder [c]	Sachets containing: - iron (elemental) 12.5 mg (as coated ferrous fumarate) - zinc (elemental) 5 mg - vitamin A 300 micrograms - with or without other micronutrients at recommended daily values
nicotinamide	Tablet: 50 mg.
pyridoxine	Tablet: 25 mg (hydrochloride).
retinol	Capsule: 50 000 IU; 100 000 IU; 200 000 IU (as palmitate). Oral oily solution: 100 000 IU/mL (as palmitate) in multidose dispenser. Tablet (sugar-coated): 10 000 IU (as palmitate). Water-miscible injection: 100 000 IU (as palmitate) in 2 mL ampoule.
riboflavin	Tablet: 5 mg.
thiamine	Tablet: 50 mg (hydrochloride).

WHO Model List of Essential Medicines – 22nd List (2021)

<i>Complementary List</i>	
calcium gluconate	Injection: 100 mg/mL in 10 mL ampoule.
28. EAR, NOSE AND THROAT MEDICINES	
acetic acid [c]	Topical: 2%, in alcohol.
☐ budesonide [c] Therapeutic alternatives to be reviewed (2023)	Nasal spray: 100 micrograms per dose.
☐ ciprofloxacin [c] Therapeutic alternatives: - ofloxacin	Solution (ear drops): 0.3% (as hydrochloride).
☐ xylometazoline  [c] Therapeutic alternatives to be reviewed (2023)	Nasal spray: 0.05%.  Not in children less than 3 months.
29. MEDICINES FOR DISEASES OF JOINTS	
29.1 Medicines used to treat gout	
allopurinol	Tablet: 100 mg.
29.2 Disease-modifying anti-rheumatic drugs (DMARDs)	
chloroquine	Tablet: 100 mg; 150 mg (as phosphate or sulfate).
<i>Complementary List</i>	
azathioprine	Tablet: 50 mg.
hydroxychloroquine	Solid oral dosage form: 200 mg (as sulfate).
methotrexate	Tablet: 2.5 mg (as sodium salt).
penicillamine	Solid oral dosage form: 250 mg.
sulfasalazine	Tablet: 500 mg.
29.3 Juvenile joint diseases	
<i>Complementary List</i>	
acetylsalicylic acid* (acute or chronic use)	Suppository: 50 mg to 150 mg. Tablet: 100 mg to 500 mg. *For use for rheumatic fever, juvenile arthritis, Kawasaki disease.
30. DENTAL PREPARATIONS	
fluoride	Paste, cream or gel: containing between 1000 and 1500 ppm fluoride (any type). In other appropriate topical formulations.
glass ionomer cement	Single-use capsules: 0.4 g powder + 0.09 mL liquid. Multi-use bottle: powder + liquid. Powder (fluoro-alumino-silicate glass) contains: 25-50% silicate, 20-40% aluminium oxide, 1-20% fluoride, 15-40% metal oxide, 0-15% phosphate, remainder are polyacrylic acid powder and metals in minimal quantities. Liquid (aqueous) contains: 7-25% polybasic carboxylic acid, 45-60% polyacrylic acid.
silver diamine fluoride	Solution: 38% w/v.

WHO Model List of Essential Medicines – 22nd List (2021)

Table 1.1: Medicines with age or weight restrictions

artesunate + pyronaridine tetraphosphate	> 5 kg
atropine	> 3 months
bedaquiline	≥ 5 years
benzyl benzoate	>2 years
betamethasone topical preparations	hydrocortisone preferred in neonates
cefazolin	> 1 month
ceftriaxone	> 41 weeks corrected gestational age
darunavir	> 3 years
delamanid	≥ 3 years (25 mg dispersible tablet) ≥ 6 years (50 mg tablet)
dihydroartemisinin + piperaquine phosphate	> 5 kg
diloxanide	>25 kg
dolutegravir	≥ 4 weeks and ≥ 3 kg (10 mg dispersible tablet) ≥ 25 kg (50 mg tablet)
doxycycline	> 8 years (except for serious infections e.g. cholera)
fluoxetine	> 8 years
ibuprofen	> 3 months (except IV form for patent ductus arteriosus)
mefloquine	> 5 kg or > 3 months
metoclopramide	Not in neonates
nevirapine	> 6 weeks
ondansetron	> 1 month
silver sulfadiazine	> 2 months
tetracaine	Not in preterm neonates
xylometazoline	> 3 months

WHO Model List of Essential Medicines – 22nd List (2021)

Table 1.2: Explanation of dosage forms

A. Principal dosage forms used in EML – oral administration

Term	Definition
Solid oral dosage form	Refers to tablets or capsules or other solid dosage forms such as 'melts' that are immediate-release preparations. It implies that there is no difference in clinical efficacy or safety between the available dosage forms, and countries should therefore choose the form(s) to be listed depending on quality and availability. The term 'solid oral dosage form' is <i>never</i> intended to allow any type of modified-release tablet.
Tablets	Refers to: • uncoated or coated (film-coated or sugar-coated) tablets that are intended to be swallowed whole; • unscored and scored*; • tablets that are intended to be chewed before being swallowed; • tablets that are intended to be dispersed or dissolved in water or another suitable liquid before being swallowed; • tablets that are intended to be crushed before being swallowed. The term 'tablet' without qualification is <i>never</i> intended to allow any type of modified-release tablet.
Tablets (qualified)	Refers to a specific type of tablet: - chewable - tablets that are intended to be chewed before being swallowed; - dispersible - tablets that are intended to be dispersed in water or another suitable liquid before being swallowed; - soluble - tablets that are intended to be dissolved in water or another suitable liquid before being swallowed; - crushable - tablets that are intended to be crushed before being swallowed; - scored - tablets bearing a break mark or marks where sub-division is intended in order to provide doses of less than one tablet; - sublingual - tablets that are intended to be placed beneath the tongue. The term 'tablet' is always qualified with an additional term (in parentheses) in entries where one of the following types of tablet is intended: gastro-resistant (such tablets may sometimes be described as enteric-coated or as delayed-release), prolonged-release or another modified-release form.

* Scored tablets may be divided for ease of swallowing, provided that dose is a whole number of tablets.

WHO Model List of Essential Medicines – 22nd List (2021)

Capsules	Refers to hard or soft capsules. The term 'capsule' without qualification is <i>never</i> intended to allow any type of modified-release capsule.
Capsules (qualified)	The term 'capsule' with qualification refers to gastro-resistant (such capsules may sometimes be described as enteric-coated or as delayed-release), prolonged-release or another modified-release form.
Granules	Preparations that are issued to patient as granules to be swallowed without further preparation, to be chewed, or to be taken in or with water or another suitable liquid. The term 'granules' without further qualification is <i>never</i> intended to allow any type of modified-release granules.
Oral powder	Preparations that are issued to patient as powder (usually as single-dose) to be taken in or with water or another suitable liquid.
Oral liquid	Liquid preparations intended to be <i>swallowed</i> i.e. oral solutions, suspensions, emulsions and oral drops, including those constituted from powders or granules, but <i>not</i> those preparations intended for <i>oromucosal administration</i> e.g. gargles and mouthwashes. Oral liquids presented as powders or granules may offer benefits in the form of better stability and lower transport costs. If more than one type of oral liquid is available on the same market (e.g. solution, suspension, granules for reconstitution), they may be interchanged and in such cases should be bioequivalent. It is preferable that oral liquids do not contain sugar and that solutions for children do not contain alcohol.

B. Principal dosage forms used in EML – parenteral administration

Term	Definition
Injection	Refers to solutions, suspensions and emulsions including those constituted from powders or concentrated solutions.
Injection (qualified)	Route of administration is indicated in parentheses where relevant.
Injection (oily)	The term 'injection' is qualified by '(oily)' in relevant entries.
Intravenous infusion	Refers to solutions and emulsions including those constituted from powders or concentrated solutions.

C. Other dosage forms

Mode of administration	Term to be used
To the eye	Eye drops, eye ointments.
Topical	For liquids: lotions, paints. For semi-solids: cream, ointment.
Rectal	Suppositories, gel or solution.
Vaginal	Pessaries or vaginal tablets.
Inhalation	Powder for inhalation, pressurized inhalation, nebulizer.

WHO Model List of Essential Medicines – 22nd List (2021)

Index

abacavir	18	bleomycin	25
abacavir + lamivudine	19	bortezomib	30
abiraterone	31	budesonide	53, 55
acetazolamide	48	budesonide + formoterol	53
acetic acid	55	bupivacaine	1
acetylcysteine	4	bupropion	52
acetylsalicylic acid	2, 24, 38, 55	caffeine citrate	50
acyclovir	18, 47	calamine	39
adalimumab	25	calcipotriol	39
albendazole	6	calcium	54
alcohol based hand rub	40	calcium folinate	25
allopurinol	32, 55	calcium gluconate	4, 55
all-trans retinoic acid (ATRA)	30	capecitabine	26
alteplase	38	carbamazepine	5, 52
amidotrizoate	40	carbetocin	50
amikacin	8, 16	carboplatin	26
amiloride	41	cefalexin	9
amiodarone	36	cefazolin	9
amitriptyline	3, 52	cefiderocol	14
amlodipine	36	cefixime	12
amodiaquine	21	cefotaxime	12
amodiaquine – sulfadoxine + pyrimethamine	23	ceftazidime	14
amoxicillin	8	ceftazidime + avibactam	14
amoxicillin + clavulanic acid	8, 16	ceftriaxone	13
amphotericin B	17, 21	cefuroxime	13
ampicillin	9	charcoal, activated	4
anastrozole	31	chlorambucil	26
anti-D immunoglobulin	34	chloramphenicol	9
anti-rabies immunoglobulin	34	chlorhexidine	40, 50
anti-rabies virus monoclonal antibodies	45	chlorine base compound	40
anti-tetanus immunoglobulin	34	chloroquine	22, 23, 55
antivenom immunoglobulin	45	chloroxylenol	41
aprepitant	42	chlorpromazine	51
arsenic trioxide	25	cholera vaccine	46
artemether	21	ciclosporin	25
artemether + lumefantrine	21	ciprofloxacin	13, 55
artesunate	22	cisplatin	26
artesunate + amodiaquine	22	clarithromycin	13
artesunate + mefloquine	22	clindamycin	10
artesunate + pyronaridine tetraphosphate	22	clofazimine	15, 16
ascorbic acid	54	clomifene	49
asparaginase	25	clomipramine	52
atazanavir + ritonavir	18	clopidogrel	38
atracurium	47	clotrimazole	17
atropine	1, 4, 48	cloxacillin	10
azathioprine	25, 55	clozapine	51
azithromycin	12, 47	coagulation factor IX	35
barium sulfate	40	coagulation factor VIII	35
BCG vaccine	46	coal tar	39
bedaquiline	16	codeine	2
bendamustine	25	colecalfiferol	54
benzathine benzylpenicillin	9	colistin	14
benznidazole	24	condoms	49
benzoyl peroxide	39	copper-containing device	49
benzyl benzoate	39	cyclizine	3
benzylpenicillin	9	cyclophosphamide	26
betamethasone	39	cyclospore	16
bevacizumab	48	cytarabine	26
bicalutamide	31	dabigatran	33
bipenden	33	dacarbazine	27
bisoprolol	35, 36, 37	daclatasvir	20

WHO Model List of Essential Medicines – 22nd List (2021)

daclatasvir + sofosbuvir	20	fluconazole	17
dactinomycin	27	flucytosine	17
dapsone	15	fludarabine	27
darunavir	18	fludrocortisone	43
dasabuvir	20	fluorescein	40
dasatinib	30	fluoride	55
daunorubicin	27	flurouracil	28, 39
deferoxamine	4, 34	floxetine	3, 52
delamanid	16	fluphenazine	51
dengue vaccine	46	folic acid	33
desmopressin	34	fomepizole	4
dexamethasone	3, 4, 31, 42, 50	fosfomycin	14
dextran 70	35	fresh-frozen plasma	34
diaphragms	49	furosemide	37, 41
diazepam	3, 5, 52	gemcitabine	28
diazoxide	44	gentamicin	10, 47
diethylcarbamazine	6	glass ionomer cement	55
digoxin	35, 37	glecaprevir + pibrentasvir	20
dihydroartemisinin + piperaquine phosphate	22	gliclazide	44
diloxanide	21	glucagon	44
dimercaprol	4	glucose	53
diphtheria antitoxin	45	glucose with sodium chloride	53
diphtheria vaccine	46	glutaral	41
docetaxel	27	glyceryl trinitrate	35
docusate sodium	3	griseofulvin	17
dolutegravir	19	Haemophilus influenzae type b vaccine	46
dolutegravir + lamivudine + tenofovir	19	haloperidol	3, 51
dopamine	38	halothane	1
doxorubicin	27	heparin sodium	33, 34
doxycycline	10, 22, 23	hepatitis A vaccine	46
efavirenz	18	hepatitis B vaccine	46
efavirenz + emtricitabine + tenofovir	19	human papilloma virus (HPV) vaccine	46
efavirenz + lamivudine + tenofovir	19	hydralazine	36
eflornithine	24	hydrochlorothiazide	36, 37, 41
empagliflozin	44	hydrocortisone	4, 31, 39, 42, 43
emtricitabine + tenofovir	19	hydroxocobalamin	33
enalapril	36, 37	hydroxycarbamide	28, 34
enoxaparin	33	hydroxychloroquine	55
entecavir	20	hyoscine butylbromide	3
ephedrine	1	hyoscine hydrobromide	3
epinephrine (adrenaline)	4, 35, 48, 53	ibrutinib	30
equine rabies immunoglobulin	45	ibuprofen	2, 24, 51
ergocalciferol	54	ifosfamide	28
ergometrine	50	imatinib	30
ertotinib	30	influenza vaccine	47
erythromycin	47	insulin injection (soluble)	44
erythropoiesis-stimulating agents	33	intermediate-acting insulin	44
estradiol cypionate + medroxyprogesterone acetate	49	Intraperitoneal dialysis solution (of appropriate composition)	51
ethambutol	15	iodine	54
ethambutol + isoniazid + pyrazinamide + rifampicin	15	iohexol	40
ethambutol + isoniazid + rifampicin	15	ipratropium bromide	53
ethanol	40	irinotecan	28
ethinylestradiol + etonogestrel	49	isoflurane	1
ethinylestradiol + levonorgestrel	49	isoniazid	15
ethinylestradiol + norethisterone	49	isoniazid + pyrazinamide + rifampicin	15
ethionamide	16	isoniazid + pyridoxine + sulfamethoxazole + trimethoprim	19
ethosuximide	6	isoniazid + rifampicin	15
etonogestrel-releasing implant	49	isoniazid + rifapentine	15
etoposide	27	isosorbide dinitrate	35
everolimus	30	itraconazole	17
fentanyl	2	ivermectin	6, 24
ferrous salt	33	Japanese encephalitis vaccine	46
ferrous salt + folic acid	33	ketamine	1
fexinidazole	23		
filgrastim	31		

WHO Model List of Essential Medicines – 22nd List (2021)

lactulose	3	natamycin	48
lamivudine	18	neostigmine	47
lamivudine + zidovudine	19	nevirapine	18
lamotrigine	5	niclosamide	6
latanoprost	48	nicotinamide	54
ledipasvir + sofosbuvir	20	nicotine replacement therapy (NRT)	52
lenalidomide	31	nifedipine	50
leuporelin	32	nifurtimox	24
levamisole	6	nilotinib	30
levodopa + carbidopa	33	nitrofurantoin	11
levofloxacin	16	nitrous oxide	1
levonorgestrel	49	nivolumab	31
levonorgestrel-releasing implant	49	norethisterone enantate	49
levonorgestrel-releasing intrauterine system	49	normal immunoglobulin	35
levothyroxine	44	nystatin	17
lidocaine	1, 35	ofloxacin	48
lidocaine + epinephrine (adrenaline)	1	ombitasvir + paritaprevir + ritonavir	20
linezolid	14, 16	omeprazole	41
lisinopril + amlodipine	36	ondansetron	3, 42
lisinopril + hydrochlorothiazide	36	oral rehydration salts	43, 53
lithium carbonate	52	oral rehydration salts – zinc sulfate	42
long-acting insulin analogues	44	oseltamivir	20
loperamide	3	oxaliplatin	28
lopinavir + ritonavir	18	oxamniquine	6
loratadine	4	oxygen	1, 2
lorazepam	5	oxytocin	50
losartan	36, 37	paclitaxel	29
Lugol's solution	45	paliperidone	51
magnesium sulfate	5	p-aminosalicylic acid	16
mannitol	41	pancreatic enzymes	41
measles vaccine	46	paracetamol	2, 24
mebendazole	6	paromomycin	21
medroxyprogesterone acetate	43, 49	pegaspargase	29
mefloquine	22, 23	pegylated interferon alfa 2a	21
meglumine antimoniate	21	penicillamine	4, 55
meclizine	40	pentamidine	23
melarsoprol	24	permethrin	39
melphalan	28	pertussis vaccine	46
meningococcal meningitis vaccine	46	phenobarbital	5
mercaptopurine	28	phenoxymethylpenicillin	11
meropenem	14, 16	phenytoin	5
meropenem + vaborbactam	14	phytomenadione	33
mesna	32	pilocarpine	48
metformin	44	piperacillin + tazobactam	14
methadone	2, 52	platelets	34
methimazole	44, 45	plazomicin	14
methotrexate	28, 55	pneumococcal vaccine	46
methyl dopa	37	podophyllum resin	39
methylprednisolone	32	poliomyelitis vaccine	46
methylthioninium chloride (methylene blue)	4	polymyxin B	14
metoclopramide	3, 42	potassium chloride	53, 54
metronidazole	11, 21	potassium ferric hexacyano-ferrate(II)-2H ₂ O (Prussian blue)	4
micafungin	17	potassium iodide	17, 44, 45
miconazole	38	potassium permanganate	39
midazolam	1, 3, 5	povidone iodine	40
mifepristone – misoprostol	50	praziquantel	6
miltefosine	21	prednisolone	4, 32, 42, 48
misoprostol	50	primaquine	22
morphine	1, 2	procaine benzylpenicillin	11
moxifloxacin	15, 16	procarbazine	29
multiple micronutrient powder	54	progesterone vaginal ring	49
multiple micronutrient supplement	50	proguanil	23
mumps vaccine	47	propofol	1
mupirocin	39	propranolol	24
naloxone	4		

WHO Model List of Essential Medicines – 22nd List (2021)

propylthiouracil	44, 45	sulfamethoxazole + trimethoprim	11, 23
prostaglandin E1	51	sulfasalazine	42, 55
protamine sulfate	33	sumatriptan	24
pyrantel	6	suramin sodium	23
pyrazinamide	15	surfactant	51
pyridostigmine	47	suxamethonium	47
pyridoxine	54	tacrolimus	25
pyrimethamine	23	tamoxifen	32
quinine	22	telmisartan + amlodipine	37
rabies vaccine	47	telmisartan + hydrochlorothiazide	37
raltegravir	19	tenofovir disoproxil fumarate	18, 20
ranitidine	41	terbinafine	38
rasburicase	32	testosterone	43
realgar-Indigo naturalis formulation	29	tetanus vaccine	46
red blood cells	34	tetracaine	48
retinol	54	tetracycline	48
ribavirin	19, 21	thalidomide	31
riboflavin	54	thiamine	54
rifabutin	15	tick-borne encephalitis vaccine	46
rifampicin	15	timolol	48
rifapentine	15	tioguanine	29
risperidone	51	tiotropium	53
ritonavir	18	tranexamic acid	34, 50
rituximab	30	trastuzumab	30
rotavirus vaccine	46	triclabendazole	6
rubella vaccine	46	trimethoprim	12
salbutamol	53	tropicamide	40
salicylic acid	39	tuberculin, purified protein derivative (PPD)	45
selenium sulfide	38	typhoid vaccine	47
senna	3, 42	ulipristal	49
silver diamine fluoride	55	urea	39
silver sulfadiazine	39	valganciclovir	19, 20
simvastatin	38	valproic acid (sodium valproate)	5, 6, 52
sodium calcium edetate	4	vancomycin	14
sodium chloride	54	varenicline	52
sodium hydrogen carbonate	54	varicella vaccine	47
sodium lactate	54	vecuronium	47
sodium nitrite	4	verapamil	35
sodium nitroprusside	37	vinblastine	29
sodium stibogluconate	21	vincristine	29
sodium thiosulfate	4, 38	vinorelbine	29
sofosbuvir	20	voriconazole	17
sofosbuvir + velpatasvir	20	warfarin	34
spectinomycin	11	water for injection	54
spironolactone	38, 41	whole blood	34
streptokinase	38	xylometazoline	55
streptomycin	16	yellow fever vaccine	46
succimer	4	zidovudine	18
sulfadiazine	23	zinc sulfate	43
sulfadoxine + pyrimethamine	22, 23	zoledronic acid	32

MUHAMMAD SAEED AWAN,
Section Officer (Admn-II).